

SOME EXAMPLES OF TORIC SASAKI-EINSTEIN MANIFOLDS

CRAIG VAN COEVERING

ABSTRACT. A series of examples of toric Sasaki-Einstein 5-manifolds is constructed which first appeared in the author's Ph.D. thesis [40]. These are submanifolds of the toric 3-Sasakian 7-manifolds of C. Boyer and K. Galicki. And there is a unique toric quasi-regular Sasaki-Einstein 5-manifold associated to every simply connected toric 3-Sasakian 7-manifold. Using 3-Sasakian reduction as in [8, 7] an infinite series of examples is constructed of each odd second Betti number. They are all diffeomorphic to $\#kM_\infty$, where $M_\infty \cong S^2 \times S^3$, for k odd. We then make use of the same framework to construct positive Ricci curvature toric Sasakian metrics on the manifolds $X_\infty \# kM_\infty$ appearing in the classification of simply connected smooth 5-manifolds due to Smale and Barden. These manifolds are not spin, thus do not admit Sasaki-Einstein metrics. They are already known to admit toric Sasakian metrics (cf. [9]) which are not of positive Ricci curvature. We then make use of the join construction of C. Boyer and K. Galicki first appearing in [6], see also [9], to construct infinitely many toric Sasaki-Einstein manifolds with arbitrarily high second Betti number of every dimension $2m + 1 \geq 5$. This is in stark contrast to the analogous case of Fano manifolds in even dimensions.

1. INTRODUCTION

A new series of quasi-regular Sasaki-Einstein 5-manifolds is constructed. These examples first appeared in the author's Ph.D. thesis [40]. They are toric, and arise as submanifolds of toric 3-Sasakian 7-manifolds. Applying 3-Sasakian reduction to torus actions on spheres C. Boyer, K. Galicki, *et al* [8] produced infinitely many toric 3-Sasakian 7-manifolds. More precisely, one has a 3-Sasakian 7-manifold $\mathcal{S}_\Omega$ for each integral weight matrix Ω satisfying some conditions to ensure smoothness. This produces infinitely many examples of each $b_2(\mathcal{S}_\Omega) \geq 1$. A result of D. Calderbank and M. Singer [12] shows that, up to finite coverings, this produces all examples of toric 3-Sasakian 7-manifolds. Associated to each $\mathcal{S}_\Omega$ is its twistor space $\mathcal{Z}$, a complex contact Fano 3-fold with orbifold singularities. The action of T^2 complexifies to $T_{\mathbb{C}}^2 = \mathbb{C}^* \times \mathbb{C}^*$ acting on $\mathcal{Z}$. Furthermore, if $\mathbf{L}$ is the line bundle of the complex contact structure, the action defines a pencil

$$E = \mathbb{P}(\mathfrak{t}_{\mathbb{C}}) \subseteq |\mathbf{L}|,$$

where $\mathfrak{t}_{\mathbb{C}}$ is the Lie algebra of $T_{\mathbb{C}}^2$. We determine the structure of the divisors in E . The generic $X \in E$ is a toric variety with orbifold structure whose orbifold anti-canonical bundle $\mathbf{K}_X^{-1}$ is positive. The total space M of the associated S^1 orbifold bundle to $\mathbf{K}_X$ has a natural Sasaki-Einstein structure. Associated to any

Date: January 18, 2007.

1991 *Mathematics Subject Classification.* Primary 53C25, Secondary 53C55, 14M25 .

Key words and phrases. Sasakian manifold, Einstein metric, toric varieties.

toric 3-Sasakian 7-manifold $\mathcal{S}$ with $\pi_1(\mathcal{S}) = e$ we have the following diagram.

$$(1.1) \quad \begin{array}{ccc} M & \longrightarrow & \mathcal{S} \\ \downarrow & & \downarrow \\ X & \longrightarrow & \mathcal{Z} \\ & & \downarrow \\ & & \mathcal{M} \end{array}$$

The horizontal maps are inclusions and the vertical are orbifold fibrations. And $\mathcal{M}$ is the 4-dimensional anti-self-dual Einstein orbifold over which $\mathcal{S}$ is an $Sp(1)$ or $SO(3)$ orbifold bundle.

It follows from the Smale/Barden classification of smooth 5-manifolds (cf. [38] and [3]) that M is diffeomorphic to $\#k(S^2 \times S^3)$ where $b_2(M) = k$. We have the following:

Theorem 1.1. *Associated to every simply connected toric 3-Sasakian 7-manifold $\mathcal{S}$ is a toric quasi-regular Sasaki-Einstein 5-manifold M . If $b_2(\mathcal{S}) = k$, then $b_2(M) = 2k + 1$ and*

$$M \underset{\text{diff}}{\cong} \#m(S^2 \times S^3), \text{ where } m = b_2(M).$$

This gives an invertible correspondence. That is, given either X or M in diagram 1.1 one can recover the other spaces with their respective geometries.

This gives an infinite series of quasi-regular Sasaki-Einstein structures on $\#m(S^2 \times S^3)$ for every odd $m \geq 3$.

In section 2 we review the basics of Sasakian geometry. In section 3 we cover the toric geometry used in the proof of 1.1. The reader can find a complete proof of the solution to the Einstein equations giving theorem 1.1 in [40]. But the existence problem of Sasaki-Einstein structures on toric Sasakian manifolds is completely solved in [41] and [18]. So this result is summarized in section 4. The basics of 3-Sasakian manifolds and 3-Sasakian reduction are covered in section 5. This section also contains some results on anti-self-dual Einstein orbifolds such as the classification result of D. Calderbank and M. Singer used in the correspondence in theorem 1.1. Section 6 contains the proof of theorem 1.1 and completes diagram 1.1. The only other possible topological types for simply connected toric Sasakian 5-manifolds are the non-spin manifolds $X_\infty \# kM_\infty$. In section 7 we use the framework already constructed to construct positive Ricci curvature Sasakian metrics on these manifolds. They are already shown to admit Sasakian structures in [9]. In section 8 we use the join construction of C. Boyer and K. Galicki [6, 9] to construct higher dimensional examples of toric quasi-regular Sasaki-Einstein and positive Ricci curvature manifolds. In particular, in every possible dimension $n \geq 5$ there exist infinitely many toric quasi-regular Sasaki-Einstein manifolds with arbitrarily high second Betti number.

This article concentrates on the quasi-regular case. This is not for lack of interest in irregular Sasaki-Einstein manifolds, but that case is covered well elsewhere, such as in [31, 32, 18].

This article makes copious use of orbifolds, V-bundles on them, and orbifold invariants such as the orbifold fundamental group π_1^{orb} and orbifold cohomology H_{orb}^* . We will use the terminology 'V-bundle' to denote an orbifold bundle. The characteristic classes of V-bundles will be elements of orbifold cohomology. The

reader unfamiliar with these concepts can consult [24] or the appendices of [5] or [40].

2. SASAKIAN MANIFOLDS

We summarize the basics of Sasakian geometry in this section. See the survey article of C. Boyer and K. Galicki [5] for more details.

Definition 2.1. *Let (M, g) be a Riemannian manifold of dimension $n = 2m + 1$, and ∇ the Levi-Civita connection. Then (M, g) is Sasakian if either of the following equivalent conditions hold:*

- (i) *There exists a unit length Killing vector field ξ on M so that the $(1, 1)$ tensor $\Phi(X) = \nabla_X \xi$ satisfies the condition*

$$(\nabla_X \Phi)(Y) = g(\xi, Y)X - g(X, Y)\xi$$

for vector fields X and Y on M .

- (ii) *The metric cone $C(M) = \mathbb{R}_+ \times M, \bar{g} = dr^2 + r^2g$ is Kähler.*

Define η to be the one form dual to ξ , i.e. $\eta(X) = g(X, \xi)$. We say that $\{g, \Phi, \xi, \eta\}$ defines a Sasakian structure on M . Note that $D = \ker \eta$ defines a contact structure on M , and Φ defines a CR -structure on D . Also, the integral curves of ξ are geodesics. The one form η extends to a one form on $C(M)$ as $\eta(X) = \frac{1}{r^2} \bar{g}(\xi, X)$. In (ii) of the definition M is identified with the subset $r = 1$ of $C(M)$, and $\xi = Jr\partial_r$. And the complex structure arises as

$$(2.1) \quad Jr\partial_r = \xi \quad JY = \Phi(Y) - \eta(Y)r\partial_r \quad \text{for } Y \in TM.$$

The Kähler form of $(C(M), \bar{g})$ is given by

$$(2.2) \quad \frac{1}{2}d(r^2\eta) = \frac{1}{2}dd^c r^2,$$

where $d^c = \frac{i}{2}(\bar{\partial} - \partial)$.

Besides the Kähler structure on $C(M)$ there is *transverse Kähler structure* on M . The Killing vector field ξ generates the Reeb foliation $\mathcal{F}_\xi$ on M . The vector field $\xi - iJ\xi = \xi + ir\partial_r$ on $C(M)$ is holomorphic and generates a local $\mathbb{C}^*$ -action extending that of ξ on M . The local orbits of this action define a transverse holomorphic structure on $\mathcal{F}_\xi$. One can choose an open covering $\{U_\alpha\}_{\alpha \in A}$ of M such that we have the projection onto the local leaf space $\pi_\alpha : U_\alpha \rightarrow V_\alpha$. Then when $U_\alpha \cap U_\beta \neq \emptyset$, the transition

$$\pi_\beta \circ \pi_\alpha^{-1} : \pi_\alpha(U_\alpha \cap U_\beta) \rightarrow \pi_\beta(U_\alpha \cap U_\beta)$$

is holomorphic. There is an isomorphism

$$d\pi_\alpha : D_p \rightarrow T_{\pi_\alpha(p)}V_\alpha,$$

for each $p \in U_\alpha$, which allows one to define a metric g^T with Kähler form $\omega = \frac{1}{2}d\eta$ as restrictions of g and $\frac{1}{2}d\eta = g(\Phi \cdot, \cdot)$ to D_p . These are easily seen to be invariant under coordinate changes. Straight forward calculation gives the following:

Proposition 2.2. *Let (M, g) be a Sasakian manifold, and $\pi_\alpha : U_\alpha \rightarrow V_\alpha$ as above. If $Y, Z \in \Gamma(D)$ are π_α -related to $\tilde{Y}, \tilde{Z} \in \Gamma(TV_\alpha)$ then*

$$(2.3) \quad \text{Ric}^T(\tilde{Y}, \tilde{Z}) = \text{Ric}(Y, Z) + 2g(X, Y),$$

$$(2.4) \quad s^T = s + 2m,$$

where Ric, s , resp. Ric^T, s^T , are the Ricci and scalar curvatures of g , resp. g^T .

Definition 2.3. A Sasaki-Einstein manifold is a Sasakian manifold (M, g) with $\text{Ric} = 2mg$.

Note that one always has $\text{Ric}(\xi, \xi) = 2m$ which fixes the Einstein constant at $2m$. Simple calculation shows that (M, g) is Einstein if, only if, $(C(M), \bar{g})$ is Ricci flat.

If ξ on M induces a free S^1 -action then $\{g, \Phi, \xi, \eta\}$ is a *regular* Sasakian structure. Another possibility is that all the orbits close but the action is not free, then the structure is *quasi-regular*. The third possibility is that not all the orbits close, in which case the generic orbit does not close. In this case the Sasakian structure is *irregular*. In the regular, resp. quasi-regular, cases the leaf space, along with its transverse Kähler structure, is a manifold, resp. orbifold, X . And M is the total space of a principal S^1 -bundle, resp. S^1 V-bundle, $\pi : M \rightarrow X$. A V-bundle is the orbifold analogue of a fiber bundle. See the appendices of [5] or [40] for details.

For a quasi-regular Sasakian manifold (M, g, Φ, ξ, η) , the leaf space X of $\mathcal{F}_\xi$ is a normal projective, algebraic variety with an orbifold structure and a Kähler form ω with $[\omega] \in H_{orb}^2(X, \mathbb{Z})$. We will make use of the following well known converse [6].

Proposition 2.4. Let (X, ω) be a Kähler orbifold, with $[\omega] \in H_{orb}^2(X, \mathbb{Z})$. There is a holomorphic line V-bundle $\mathbf{L}$ with $c_1(\mathbf{L}) = -[\omega] \in H_{orb}^2(X, \mathbb{Z})$ and an S^1 -principal subbundle $M \subset \mathbf{L}$ such that M has a family of Sasakian structures $\{g_a, \Phi, \xi_a, \eta_a\}, a \in \mathbb{R}_+$,

$$g_a = a^2 \eta \otimes \eta + a\pi^* h, \text{ where } h \text{ is a Kähler metric on } X.$$

Furthermore, if $\mathbf{L}^\times$ is $\mathbf{L}$ minus the zero section, then $\mathbf{L}^\times$ is biholomorphic to $C(M)$.

If (X, ω) is Kähler-Einstein with positive scalar curvature, and $q[\omega] \in c_1^{orb}(X)$, for $q \in \mathbb{Z}_+$, then exactly one of the above Sasakian structures is Sasaki-Einstein, and one may take $\mathbf{L} = K_X^{\frac{1}{q}}$.

Note that if $\pi_1^{orb}(X) = e$ and $[\omega]$ is indivisible in $H_{orb}^2(X, \mathbb{Z})$, then $\pi_1^{orb}(M) = e$. Of course we are interested in the case where M is smooth. This happens when the action of the local uniformizing groups of X inject into the group of the fibers of the V-bundle M .

We will mainly be interested in *toric* Sasaki-Einstein manifolds.

Definition 2.5. A toric Sasakian manifold is a Sasakian manifold M of dimension $2m + 1$ whose Sasakian structure $\{g, \Phi, \xi, \eta\}$ is preserved by an effective action of an $(m + 1)$ -dimensional torus T such that ξ is an element of the Lie algebra $\mathfrak{t}$ of T .

Let $T_{\mathbb{C}} \cong (\mathbb{C}^*)^{m+1}$ be the complexification of T , then $T_{\mathbb{C}}$ acts on $C(M)$ by holomorphic automorphisms. One sees that this definition is equivalent to $C(M)$ being a toric Kähler manifold.

3. TORIC GEOMETRY

We give some basic definitions in the theory of toric varieties that we will need. See [16, 33, 34] for more details. In addition we will consider the notion of a compatible orbifold structure on a toric variety and holomorphic V-bundles. We are interested in Kähler toric orbifolds, and will give a description of the Kähler structure due to V. Guillemin [22].

3.1. Toric varieties. Let $N \cong \mathbb{Z}^r$ be the free $\mathbb{Z}$ -module of rank r and $M = \text{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$ its dual. We denote $N_{\mathbb{Q}} = N \otimes \mathbb{Q}$ and $M_{\mathbb{Q}} = M \otimes \mathbb{Q}$ with the natural pairing

$$\langle \cdot, \cdot \rangle : M_{\mathbb{Q}} \times N_{\mathbb{Q}} \rightarrow \mathbb{Q}.$$

Similarly we denote $N_{\mathbb{R}} = N \otimes \mathbb{R}$ and $M_{\mathbb{R}} = M \otimes \mathbb{R}$.

Let $T_{\mathbb{C}} := N \otimes_{\mathbb{Z}} \mathbb{C}^* \cong \mathbb{C}^* \times \cdots \times \mathbb{C}^*$ be the algebraic torus. Each $m \in M$ defines a character $\chi^m : T_{\mathbb{C}} \rightarrow \mathbb{C}^*$ and each $n \in N$ defines a one-parameter subgroup $\lambda_n : \mathbb{C}^* \rightarrow T_{\mathbb{C}}$. In fact, this gives an isomorphism between M (resp. N) and the multiplicative group $\text{Hom}_{\text{alg.}}(T_{\mathbb{C}}, \mathbb{C}^*)$ (resp. $\text{Hom}_{\text{alg.}}(\mathbb{C}^*, T_{\mathbb{C}})$).

Definition 3.1. A subset σ of $N_{\mathbb{R}}$ is a strongly convex rational polyhedral cone if there are $n_1, \dots, n_r$ so that

$$\sigma = \mathbb{R}_{\geq 0}n_1 + \cdots + \mathbb{R}_{\geq 0}n_r,$$

and one has $\sigma \cap -\sigma = \{o\}$, where $o \in N$ is the origin.

The dimension $\dim \sigma$ is the dimension of the $\mathbb{R}$ -subspace $\sigma + (-\sigma)$ of $N_{\mathbb{R}}$. The dual cone to σ is

$$\sigma^{\vee} = \{x \in M_{\mathbb{R}} : \langle x, y \rangle \geq 0 \text{ for all } y \in \sigma\},$$

which is also a convex rational polyhedral cone. A subset τ of σ is a face, $\tau < \sigma$, if

$$\tau = \sigma \cap m^{\perp} = \{y \in \sigma : \langle m, y \rangle = 0\} \text{ for } m \in \sigma^{\vee}.$$

And τ is a strongly convex rational polyhedral cone.

Definition 3.2. A fan in N is a collection Δ of strongly convex rational polyhedral cones such that:

- (i) For $\sigma \in \Delta$ every face of σ is contained in Δ .
- (ii) For any $\sigma, \tau \in \Delta$, the intersection $\sigma \cap \tau$ is a face of both σ and τ .

We will consider *complete* fans for which the support $\bigcup_{\sigma \in \Delta} \sigma$ is $N_{\mathbb{R}}$. We will denote

$$\Delta(i) := \{\sigma \in \Delta : \dim \sigma = i\}, \quad 0 \leq i \leq n.$$

Definition 3.3. A fan in N is nonsingular if each $\sigma \in \Delta(r)$ is generate by r elements of N which can be completed to a $\mathbb{Z}$ -basis of N . A fan in N is simplicial if each $\sigma \in \Delta(r)$ is generated by r elements of N which can be completed to a $\mathbb{Q}$ -basis of $N_{\mathbb{Q}}$.

If σ is a strongly convex rational polyhedral cone, $S_{\sigma} = \sigma^{\vee} \cap M$ is a finitely generated semigroup. We denote by $\mathbb{C}[S_{\sigma}]$ the semigroup algebra. We will denote the generators of $\mathbb{C}[S_{\sigma}]$ by x^m for $m \in S_{\sigma}$. Then $U_{\sigma} := \text{Spec } \mathbb{C}[S_{\sigma}]$ is a normal affine variety on which $T_{\mathbb{C}}$ acts algebraically with a (Zariski) open orbit isomorphic to $T_{\mathbb{C}}$. If σ is nonsingular, then $U_{\sigma} \cong \mathbb{C}^n$.

Theorem 3.4 ([16, 33, 34]). For a fan Δ in N the affine varieties U_{σ} for $\sigma \in \Delta$ glue together to form an irreducible normal algebraic variety

$$X_{\Delta} = \bigcup_{\sigma \in \Delta} U_{\sigma}.$$

Furthermore, X_{Δ} is non-singular if, and only if, Δ is nonsingular. And X_{Δ} is compact if, and only if, Δ is complete.

Proposition 3.5. *The variety X_Δ has an algebraic action of $T_{\mathbb{C}}$ with the following properties.*

- (i) *To each $\sigma \in \Delta(i), 0 \leq i \leq n$, there corresponds a unique $(n-i)$ -dimensional $T_{\mathbb{C}}$ -orbit $\text{Orb}(\sigma)$ so that X_Δ decomposes into the disjoint union*

$$X_\Delta = \bigcup_{\sigma \in \Delta} \text{Orb}(\sigma),$$

where $\text{Orb}(o)$ is the unique n -dimensional orbit and is isomorphic to $T_{\mathbb{C}}$.

- (ii) *The closure $V(\sigma)$ of $\text{Orb}(\sigma)$ in X_Δ is an irreducible $(n-i)$ -dimensional $T_{\mathbb{C}}$ -stable subvariety and*

$$V(\sigma) = \bigcup_{\tau \geq \sigma} \text{Orb}(\tau).$$

We will consider toric varieties with an orbifold structure.

Definition 3.6. *We will denote by Δ^* an augmented fan by which we mean a fan Δ with elements $n(\rho) \in N \cap \rho$ for every $\rho \in \Delta(1)$.*

Proposition 3.7. *For a complete simplicial augmented fan Δ^* we have a natural orbifold structure compatible with the action of $T_{\mathbb{C}}$ on X_Δ . We denote X_Δ with this orbifold structure by X_{Δ^*} .*

Proof. Let $\sigma \in \Delta^*(n)$ have generators $p_1, p_2, \dots, p_n$ as in the definition. Let $N' \subseteq N$ be the sublattice $N' = \mathbb{Z}\{p_1, p_2, \dots, p_n\}$, and σ' the equivalent cone in N' . Denote by M' the dual lattice of N' and $T'_{\mathbb{C}}$ the torus. Then $U_{\sigma'} \cong \mathbb{C}^n$. It is easy to see that

$$N/N' = \text{Hom}_{\mathbb{Z}}(M'/M, \mathbb{C}^*).$$

And N/N' is the kernel of the homomorphism

$$T'_{\mathbb{C}} = \text{Hom}_{\mathbb{Z}}(M', \mathbb{C}^*) \rightarrow T_{\mathbb{C}} = \text{Hom}_{\mathbb{Z}}(M, \mathbb{C}^*).$$

Let $\Gamma = N/N'$. An element $t \in \Gamma$ is a homomorphism $t : M' \rightarrow \mathbb{C}^*$ equal to 1 on M . The regular functions on $U_{\sigma'}$ consist of $\mathbb{C}$ -linear combinations of x^m for $m \in \sigma'^{\vee} \cap M'$. And $t \cdot x^m = t(m)x^m$. Thus the invariant functions are the $\mathbb{C}$ -linear combinations of x^m for $m \in \sigma^{\vee} \cap M$, the regular functions of U_σ . Thus $U_{\sigma'}/\Gamma = U_\sigma$. And the charts are easily seen to be compatible on intersections. $\square$

Conversely, we have the following.

Proposition 3.8. *Let Δ be a complete simplicial fan. Suppose for simplicity that the local uniformizing groups are abelian. Then every orbifold structure on X_Δ compatible with the action of $T_{\mathbb{C}}$ arises from an augmented fan Δ^* .*

See [40] for a proof.

Let Δ^* be an augmented fan in N . We will assumed from now on that the fan Δ is simplicial and complete.

Definition 3.9. *A real function $h : N_{\mathbb{R}} \rightarrow \mathbb{R}$ is a Δ^* -linear support function if for each $\sigma \in \Delta^*$ with given $\mathbb{Q}$ -generators $p_1, \dots, p_r$ in N , there is an $l_\sigma \in M_{\mathbb{Q}}$ with $h(s) = \langle l_\sigma, s \rangle$ and l_σ is $\mathbb{Z}$ -valued on the sublattice $\mathbb{Z}\{p_1, \dots, p_r\}$. And we require that $\langle l_\sigma, s \rangle = \langle l_\tau, s \rangle$ whenever $s \in \sigma \cap \tau$. The additive group of Δ^* -linear support functions will be denoted by $\text{SF}(\Delta^*)$.*

Note that $h \in \text{SF}(\Delta^*)$ is completely determined by the integers $h(n(\rho))$ for all $\rho \in \Delta(1)$. And conversely, an assignment of an integer to $h(n(\rho))$ for all $\rho \in \Delta(1)$ defines h . Thus

$$\text{SF}(\Delta^*) \cong \mathbb{Z}^{\Delta(1)}.$$

Definition 3.10. Let Δ^* be a complete augmented fan. For $h \in \text{SF}(\Delta^*)$,

$$\Sigma_h := \{m \in M_{\mathbb{R}} : \langle m, n \rangle \geq h(n), \text{ for all } n \in N_{\mathbb{R}}\},$$

is a, possibly empty, convex polytope in $M_{\mathbb{R}}$.

We will consider the holomorphic line V -bundles on $X = X_{\Delta^*}$. All V -bundles will be *proper* in this section. The set of isomorphism classes of holomorphic line V -bundles is denoted by $\text{Pic}^{\text{orb}}(X)$, which is a group under the tensor product.

Definition 3.11. A Baily divisor is a $\mathbb{Q}$ -Weil divisor $D \in \text{Weil}(X) \otimes \mathbb{Q}$ whose inverse image $D_{\tilde{U}} \in \text{Weil}(\tilde{U})$ in every local uniformizing chart $\pi : \tilde{U} \rightarrow U$ is Cartier. The additive group of Baily divisors is denoted $\text{Div}^{\text{orb}}(X)$.

A Baily divisor D defines a holomorphic line V -bundle $[D] \in \text{Pic}^{\text{orb}}(X)$ in a way completely analogous to Cartier divisors. Given a nonzero meromorphic function $f \in \mathcal{M}$ we have the *principal divisor*

$$\text{div}(f) := \sum_V \nu_V(f) V,$$

where $\nu_V(f)V$ is the order of the zero, or negative the order of the pole, of f along each irreducible subvariety of codimension one. We have the exact sequence

$$(3.1) \quad 1 \rightarrow \mathbb{C}^* \rightarrow \mathcal{M}^* \rightarrow \text{Div}^{\text{orb}}(X) \xrightarrow{[\cdot]} \text{Pic}^{\text{orb}}(X).$$

A holomorphic line V -bundle $\pi : \mathbf{L} \rightarrow X$ is equivariant if there is an action of $T_{\mathbb{C}}$ on $\mathbf{L}$ such that π is equivariant, $\pi(tw) = t\pi(w)$ for $w \in \mathbf{L}$ and $t \in T_{\mathbb{C}}$ and the action lifts to a holomorphic action, linear on the fibers, over each uniformizing neighborhood. The group of isomorphism classes of equivariant holomorphic line V -bundles is denoted $\text{Pic}^{\text{orb}}_{T_{\mathbb{C}}}(X)$. Similarly, we have invariant Baily divisors, denoted $\text{Div}^{\text{orb}}_{T_{\mathbb{C}}}(X)$, and $[D] \in \text{Pic}^{\text{orb}}_{T_{\mathbb{C}}}(X)$ whenever $D \in \text{Div}^{\text{orb}}_{T_{\mathbb{C}}}(X)$.

Proposition 3.12. Let $X = X_{\Delta^*}$ be compact with the standard orbifold structure, i.e. Δ^* is simplicial and complete.

- (i) There is an isomorphism $\text{SF}(\Delta^*) \cong \text{Div}^{\text{orb}}_{T_{\mathbb{C}}}(X)$ obtained by sending $h \in \text{SF}(\Delta^*)$ to

$$D_h := - \sum_{\rho \in \Delta(1)} h(n(\rho)) V(\rho).$$

- (ii) There is a natural homomorphism $\text{SF}(\Delta^*) \rightarrow \text{Pic}^{\text{orb}}_{T_{\mathbb{C}}}(X)$ which associates an equivariant line V -bundle $\mathbf{L}_h$ to each $h \in \text{SF}(\Delta^*)$.
(iii) Suppose $h \in \text{SF}(\Delta^*)$ and $m \in M$ satisfies

$$\langle m, n \rangle \geq h(n) \text{ for all } n \in N_{\mathbb{R}},$$

then m defines a section $\psi : X \rightarrow \mathbf{L}_h$ which has the equivariance property $\psi(tx) = \chi^m(t)(t\psi(x))$.

- (iv) The set of sections $H^0(X, \mathcal{O}(\mathbf{L}_h))$ is the finite dimensional $\mathbb{C}$ -vector space with basis $\{x^m : m \in \Sigma_h \cap M\}$.

- (v) Every Baily divisor is linearly equivalent to a $T_{\mathbb{C}}$ -invariant Baily divisor. Thus for $D \in \text{Pic}^{\text{orb}}(X)$, $[D] \cong [D_h]$ for some $h \in \text{SF}(\Delta^*)$.
- (vi) If $\mathbf{L}$ is any holomorphic line V -bundle, then $\mathbf{L} \cong \mathbf{L}_h$ for some $h \in \text{SF}(\Delta^*)$. The homomorphism in part i. induces an isomorphism $\text{SF}(\Delta^*) \cong \text{Pic}^{\text{orb}}_{T_{\mathbb{C}}}(X)$ and we have the exact sequence

$$0 \rightarrow M \rightarrow \text{SF}(\Delta^*) \rightarrow \text{Pic}^{\text{orb}}(X) \rightarrow 1.$$

Proof. (i) For each $\sigma \in \Delta(n)$ with uniformizing neighborhood $\pi : U_{\sigma'} \rightarrow U_{\sigma}$ above the map $h \rightarrow D_h$ assigns the principal divisor

$$\text{div}(x^{-l_{\sigma}}) = - \sum_{\rho \in \Delta(1), \rho < \sigma} h(n(\rho))V'(\rho),$$

where $V'(\rho)$ is the closure of the orbit $\text{Orb}(\rho)$ in $U_{\sigma'}$. An element $\text{Div}^{\text{orb}}_{T_{\mathbb{C}}}(X)$ must be a sum of closures of codimension one orbits $V(\rho)$ in Proposition 3.5, and by above remarks the map is an isomorphism.

(ii) One defines $\mathbf{L}_h := [D_h]$, where $[D_h]$ is constructed as follows. Consider a uniformizing chart $\pi : U_{\sigma'} \rightarrow U_{\sigma}$ as in Proposition 3.7. Define $\mathbf{L}_h|_{U_{\sigma'}}$ to be the invertible sheaf $\mathcal{O}_{U_{\sigma'}}(D_h)$, with D_h defined on $U_{\sigma'}$ by $x^{-l_{\sigma}}$. So $\mathbf{L}_h|_{U_{\sigma'}} \cong U_{\sigma'} \times \mathbb{C}$ with an action of $T'_{\mathbb{C}}$,

$$t(x, v) = (tx, \chi^{-l_{\sigma}}(t)v) \text{ where } t \in T'_{\mathbb{C}}, (x, v) \in U_{\sigma'} \times \mathbb{C}.$$

Then $\mathbf{L}_h|_{U_{\sigma}}$ is the quotient by the subgroup $N/N' \subset T'_{\mathbb{C}}$, so it has an action of $T_{\mathbb{C}}$. And the $\mathbf{L}_h|_{U_{\sigma}}$ glue together equivariantly with respect to the action.

(iii) For $\sigma \in \Delta$ we have $\langle m, n \rangle \geq \langle l_{\sigma}, n \rangle$ for all $n \in \sigma$. Then $m - l_{\sigma} \in M' \cap \sigma'^{\vee}$ and $x^{m-l_{\sigma}}$ is a section of the invertible sheaf $\mathcal{O}_{U_{\sigma'}}(D_h)$ and is equivariant with respect to N/N' so it defines a section of $\mathbf{L}_h|_{U_{\sigma}}$. And these sections are compatible.

(iv) We will make use of the GAGA theorems of A. Grothendieck [20, 21]. As with any holomorphic V -bundle, the sheaf of sections $\mathcal{O}(\mathbf{L}_h)$ is a coherent sheaf. It follows from GAGA that we may consider $\mathcal{O}(\mathbf{L}_h)$ as a coherent algebraic sheaf, and all global sections are algebraic. If ϕ is a global section, then $\phi \in H^0(T_{\mathbb{C}}, \mathcal{O}(\mathbf{L}_h)) \subset \mathbb{C}[M]$. And in the uniformizing chart $\pi : U_{\sigma'} \rightarrow U_{\sigma}$, ϕ lifts to an element of the module $\mathcal{O}_{U_{\sigma'}} \cdot x^{l_{\sigma}}$ which has a basis $\{x^m : m \in l_{\sigma} + M' \cap \sigma'^{\vee}\}$. So $\phi|_{U_{\sigma}}$ is a $\mathbb{C}$ -linear combination of x^m with $m \in M$ and $\langle m, n \rangle \geq h(n)$ for all $n \in \sigma$. Thus $m \in \Sigma_h$.

(v) The divisor $T_{\mathbb{C}} \cap D$ is a Cartier divisor on $T_{\mathbb{C}}$ which is also principal since $\mathbb{C}[M]$ is a unique factorization domain. Thus there is a nonzero rational function f so that $D' = D - \text{div}(f)$ satisfies $D' \cap T_{\mathbb{C}} = \emptyset$. Then $D' \in \text{Div}^{\text{orb}}_{T_{\mathbb{C}}}(X)$, and the result follows from i.

(vi) Consider $\mathbf{L}_{U_{\sigma'}}$ on a uniformizing neighborhood $U_{\sigma'}$ as above. For each $\rho \in \Delta(1), \rho < \sigma$ the subgroup $H_{\rho} \subseteq N/N'$ fixing $V'(\rho)$ is cyclic and generated by $n' \in N$ where n' is the primitive element with $a_{\rho}n' = n(\rho)$. Now H_{ρ} acts linearly on the fibers of $\mathbf{L}_{U_{\sigma'}}$ over $V'(\rho)$. Suppose n' acts with weight $e^{2\pi i \frac{k}{a}}$, then let $D_{\rho} := kV(\rho)$. If $D' := \sum_{\rho \in \Delta(1)} D_{\rho}$, then $\mathbf{L}' := \mathbf{L} \otimes [-D']$ is Cartier on $X_0 := X \setminus \text{Sing}(X)$, where $\text{Sing}(X)$ has codimension at least two. The sheaf $\mathcal{O}(\mathbf{L}')$ is not only coherent but is a rank-1 reflexive sheaf. By GAGA $\mathcal{O}(\mathbf{L}') \cong E \otimes \mathcal{O}'$, where E is an algebraic reflexive rank-1 sheaf and $\mathcal{O}'$ is the sheaf of analytic functions. It is well known that $E = \mathcal{O}(D)$ for $D \in \text{Weil}(X)$. And as a Baily divisor, we have $\mathbf{L}' \cong [D]$. So $\mathbf{L} \cong [D + D']$, and by v. we have $\mathbf{L} \cong \mathbf{L}_h$ for some $h \in \text{SF}(\Delta^*)$. $\square$

The sign convention in the proposition is adopted to make subsequent discussions involving Σ_h consistent with the existing literature, although having $D_{-m} = \text{div}(x^m)$ maybe bothersome. Note also that we denote a Baily divisor by a formal $\mathbb{Z}$ -linear sum the coefficient giving the multiplicity of the irreducible component in the *uniformizing chart*. This is different from its expression as a Weil divisor when irreducible components are contained in codimension-1 components of the singular set of the orbifold.

For $X = X_{\Delta^*}$ there is a unique $k \in \text{SF}(\Delta^*)$ such that $k(n(\rho)) = 1$ for all $\rho \in \Delta(1)$. The corresponding Baily divisor

$$(3.2) \quad D_k := - \sum_{\rho \in \Delta(1)} V(\rho)$$

is the (*orbifold*) *canonical divisor*. The corresponding V -bundle is $\mathbf{K}_X$, the V -bundle of holomorphic n -forms. This will in general be different from the canonical sheaf in the algebraic geometric sense.

Definition 3.13. Consider support functions as above but which are only required to be $\mathbb{Q}$ -valued on $N_{\mathbb{Q}}$, denoted $\text{SF}(\Delta, \mathbb{Q})$. h is strictly upper convex if $h(n + n') \geq h(n) + h(n')$ for all $n, n' \in N_{\mathbb{Q}}$ and for any two $\sigma, \sigma' \in \Delta(n)$, l_{σ} and $l_{\sigma'}$ are different linear functions.

Given a strictly upper convex support function h , the polytope Σ_h is the convex hull in $M_{\mathbb{R}}$ of the vertices $\{l_{\sigma} : \sigma \in \Delta(n)\}$. Each $\rho \in \Delta(1)$ defines a facet by

$$\langle m, n(\rho) \rangle \geq h(n(\rho)).$$

If $n(\rho) = a_{\rho} n'$ with $n' \in N$ primitive and $a_{\rho} \in \mathbb{Z}^+$ we may label the face with a_{ρ} to get the labeled polytope Σ_h^* which encodes the orbifold structure. Conversely, from a rational convex polytope Σ^* we associate a fan Δ^* and a support function h as follows. For an l -dimensional face $\theta \subset \Sigma^*$, define the rational n -dimensional cone $\sigma^{\vee}(\theta) \subset M_{\mathbb{R}}$ consisting of all vectors $\lambda(p - p')$, where $\lambda \in \mathbb{R}_{\geq 0}$, $p \in \Sigma$, and $p' \in \theta$. Then $\sigma(\theta) \subset N_{\mathbb{R}}$ is the $(n - l)$ -dimensional cone dual to $\sigma^{\vee}(\theta)$. The set of all $\sigma(\theta)$ defines the complete fan Δ^* , where one assigns $n(\rho)$ to $\rho \in \Delta(1)$ if $n(\rho) = an'$ with n' primitive and a is the label on the corresponding $(n - 1)$ -dimensional face of Σ^* . The corresponding rational support function is then

$$h(n) = \inf\{\langle m, n \rangle : m \in \Sigma^*\} \text{ for } n \in N_{\mathbb{R}}.$$

Proposition 3.14 ([34, 16]). *There is a one-to-one correspondence between the set of pairs (Δ^*, h) with $h \in \text{SF}(\Delta, \mathbb{Q})$ strictly upper convex, and rational convex marked polytopes Σ_h^* .*

We will be interested in toric orbifolds X_{Δ^*} with such a support function and polytope, Σ_h^* . More precisely we will be concerned with the following.

Definition 3.15. Let $X = X_{\Delta^*}$ be a compact toric orbifold. We say that X is Fano if $-k \in \text{SF}(\Delta^*)$, which defines the anti-canonical V -bundle $\mathbf{K}_X^{-1}$, is strictly upper convex.

These toric varieties are not necessarily Fano in the usual sense, since $\mathbf{K}_X^{-1}$ is the *orbifold* anti-canonical class. This condition is equivalent to $\{n \in N_{\mathbb{R}} : k(n) \leq 1\} \subset N_{\mathbb{R}}$ being a convex polytope with vertices $n(\rho), \rho \in \Delta(1)$. We will use Δ^* to denote both the augmented fan and this polytope in this case.

If $\mathbf{L}_h$ is a line V -bundle, then for certain $s > 0$, $\mathbf{L}_h^s \cong \mathbf{L}_{sh}$ will be a holomorphic line bundle. For example $s = \text{Ord}(X)$, the least common multiple of the orders of the uniformizing groups, will do. So suppose $\mathbf{L}_h$ is a holomorphic line bundle. If the global holomorphic sections generate $\mathbf{L}_h$, by Proposition 3.12 $M \cap \Sigma_h = \{m_0, m_1, \dots, m_r\}$ and we have a holomorphic map $\psi_h : X \rightarrow \mathbb{C}P^r$ where

$$(3.3) \quad \psi_h(w) := [x^{m_0}(w) : x^{m_1}(w) : \dots : x^{m_r}(w)].$$

Proposition 3.16 ([34]). *Suppose $\mathbf{L}_h$ is a line bundle, so $h \in \text{SF}(\Delta^*)$ is integral, and suppose h is strictly upper convex. Then $\mathbf{L}_h$ is ample, meaning that for large enough $\nu > 0$*

$$\psi_{\nu h} : X \rightarrow \mathbb{C}P^N,$$

is an embedding, where $M \cap \Sigma_{\nu h} = \{m_0, m_1, \dots, m_N\}$.

Corollary 3.17. *Let X be a Fano toric orbifold. If $\nu > 0$ is sufficiently large with $-\nu k$ integral, $\mathbf{K}^{-\nu}$ will be very ample and $\psi_{-\nu k} : X \rightarrow \mathbb{C}P^N$ an embedding.*

Let X_{Δ^*} be an orbifold surface with $h \in \text{SF}(\Delta^*)$. Then the total spaces of $\mathbf{L}_h$ and $\mathbf{L}_h^\times$ are toric varieties. The fan of $\mathbf{L}_h^\times$ is as follows. If $\sigma \in \Delta^*$ is spanned by $n(\rho_1), \dots, n(\rho_k) \in \mathbb{Z}^n$ as in definition 3.6 let $\bar{\sigma}$ be the cone in $\mathbb{R}^{n+1}$ spanned by $(n(\rho_1), h(n(\rho_1))), \dots, (n(\rho_k), h(n(\rho_k))) \in \mathbb{Z}^{n+1}$. The collection of $\bar{\sigma}, \sigma \in \Delta^*$ defines a fan $\mathcal{C} = \mathcal{C}_h$, which is the fan of $\mathbf{L}_h^\times$. Furthermore, if h , or $-h$, is strictly upper convex, then one can add an additional $(n+1)$ -cone to $\mathcal{C}_h$ to get an affine variety $Y = \mathbf{L}_h^\times \cup \{p\}$. We will make use of the smoothness condition on $\mathbf{L}_h^\times$. The toric variety $\mathbf{L}_h^\times$ is smooth if for every n -cone $\bar{\sigma}$ of $\mathcal{C}$ as above spanned by $\tau_1, \dots, \tau_n \in \mathbb{Z}^{n+1}$ we have

$$(3.4) \quad (\mathbb{R}_{\geq 0}\tau_1 + \dots + \mathbb{R}_{\geq 0}\tau_n) \cap \mathbb{Z}^{n+1} = \mathbb{Z}_{\geq 0}\tau_1 + \dots + \mathbb{Z}_{\geq 0}\tau_n.$$

Suppose $\mathbf{L}_h^q \cong \mathbf{K}_X$ for some $q \in \mathbb{Z}_{>0}$. Then $\mathbf{K}_Y$ is trivial. That is, Y has a Gorenstein singularity at the apex.

3.2. Kähler structures. We review the construction of toric Kähler metrics on toric varieties. Any compact toric orbifold associated to a polytope admits a Kähler metric (see [29]). Due to T. Delzant [13] and E. Lerman and S. Tolman [29] in the orbifold case, the symplectic structure is uniquely determined up to symplectomorphism by the polytope, which is the image of the moment map. This polytope is Σ_h^* of the previous section with h generalized to be real valued. There are infinitely many Kähler structures on a toric orbifold with fixed polytope Σ_h^* , but there is a canonical Kähler metric obtained by reduction. V. Guillemin gave an explicit formula [22, 11] for this Kähler metric. In particular, we show that every toric Fano orbifold admits a Kähler metric $\omega \in c_1(X)$.

Let Σ^* be a convex polytope in $M_{\mathbb{R}} \cong \mathbb{R}^{n*}$ defined by the inequalities

$$(3.5) \quad \langle x, u_i \rangle \geq \lambda_i, \quad i = 1, \dots, d,$$

where $u_i \in N \subset N_{\mathbb{R}} \cong \mathbb{R}^n$ and $\lambda_i \in \mathbb{R}$. If Σ_h^* is associated to (Δ^*, h) , then the u_i and λ_i are the set of pairs $n(\rho)$ and $h(n(\rho))$ for $\rho \in \Delta(1)$. We allow the λ_i to be real but require any set $u_{i_1}, \dots, u_{i_n}$ corresponding to a vertex to form a $\mathbb{Q}$ -basis of $N_{\mathbb{Q}}$.

Let $(e_1, \dots, e_d)$ be the standard basis of $\mathbb{R}^d$ and $\beta : \mathbb{R}^d \rightarrow \mathbb{R}^n$ be the map which takes e_i to u_i . Let $\mathfrak{n}$ be the kernel of β , so we have the exact sequence

$$(3.6) \quad 0 \rightarrow \mathfrak{n} \xrightarrow{\iota} \mathbb{R}^d \xrightarrow{\beta} \mathbb{R}^n \rightarrow 0,$$

and the dual exact sequence

$$(3.7) \quad 0 \rightarrow \mathbb{R}^{n*} \xrightarrow{\beta^*} \mathbb{R}^{d*} \xrightarrow{\iota^*} \mathfrak{n}^* \rightarrow 0.$$

Since (3.6) induces an exact sequence of lattices, we have an exact sequence

$$(3.8) \quad 1 \rightarrow N \rightarrow T^d \rightarrow T^n \rightarrow 1,$$

where the connected component of the identity of N is an $(d-n)$ -dimensional torus. The standard representation of T^d on $\mathbb{C}^d$ preserves the Kähler form

$$(3.9) \quad \frac{i}{2} \sum_{k=1}^d dz_k \wedge d\bar{z}_k,$$

and is Hamiltonian with moment map

$$(3.10) \quad \mu(z) = \frac{1}{2} \sum_{k=1}^d |z_k|^2 e_k + c,$$

unique up to a constant c . We will set $c = \sum_{k=1}^d \lambda_k e_k$. Restricting to $\mathfrak{n}^*$ we get the moment map for the action of N on $\mathbb{C}^d$

$$(3.11) \quad \mu_N(z) = \frac{1}{2} \sum_{k=1}^d |z_k|^2 \alpha_k + \lambda,$$

with $\alpha_k = \iota^* e_k$ and $\lambda = \sum \lambda_k \alpha_k$. Let $Z = \mu_N^{-1}(0)$ be the zero set. By the exactness of (3.7) $z \in \mu_N^{-1}(0)$ if and only if there is a $v \in \mathbb{R}^{n*}$ with $\mu(z) = \beta^* v$. Since β^* is injective, we have a map

$$(3.12) \quad \nu : Z \rightarrow \mathbb{R}^{n*},$$

where $\beta^* \nu(z) = \mu(z)$ for all $z \in Z$. For $z \in Z$

$$(3.13) \quad \begin{aligned} \langle \nu(z), u_i \rangle &= \langle \beta^* \nu(z), e_i \rangle \\ &= \langle \mu(z), e_i \rangle \\ &= \frac{1}{2} |z_i|^2 + \lambda_i, \end{aligned}$$

thus $\nu(z) \in \Sigma^*$. Conversely, if $v \in \Sigma^*$, then $v = \nu(z)$ for some $z \in Z$ and in fact a T^d orbit in Z . Thus Z is compact. The following is not difficult to show.

Theorem 3.18. *The action of N on Z is locally free. Thus the quotient*

$$X_{\Sigma^*} = Z/N$$

is a compact orbifold. Let

$$\pi : Z \rightarrow X$$

be the projection and

$$\iota : Z \rightarrow \mathbb{C}^d$$

the inclusion. Then X_{Σ^} has a canonical Kähler structure with Kähler form ω uniquely defined by*

$$\pi^* \omega = \iota^* \left(\frac{i}{2} \sum_{k=1}^d dz_k \wedge d\bar{z}_k \right).$$

We have an action of $T^n = T^d/N$ on X_{Σ^*} which is Hamiltonian for ω . The map ν is T^d invariant, and it descends to a map, which we also call ν ,

$$(3.14) \quad \nu : X_{\Sigma^*} \rightarrow \mathbb{R}^{n^*},$$

which is the moment map for this action. The above comments show that $\text{Im}(\nu) = \Sigma^*$. The action T^n extends to the complex torus $T_{\mathbb{C}}^n$ and one can show that as an analytic variety and orbifold X_{Σ^*} is the toric variety constructed from Σ^* in the previous section. See [23] for more details.

Let $\sigma : \mathbb{C}^d \rightarrow \mathbb{C}^d$ be the involution $\sigma(z) = \bar{z}$. The set Z is stable under σ , and σ descends to an involution on X . We denote the fixed point sets by Z_r and X_r . And we have the projection

$$(3.15) \quad \pi : Z_r \rightarrow X_r.$$

We equip Z_r and X_r with Riemannian metrics by restricting the Kähler metrics on $\mathbb{C}^d$ and X respectively.

Proposition 3.19. *The map (3.15) is a locally finite covering and is an isometry with respect to these metrics*

Note that Z_r is a subset of $\mathbb{R}^d$ defined by

$$(3.16) \quad \frac{1}{2} \sum_{k=1}^d x_k^2 \alpha_k = -\lambda.$$

Restrict to the orthant $x_k > 0$ $k = 1, \dots, d$ of $\mathbb{R}^d$. Let Z'_r be the component of Z_r in this orthant. Under the coordinates

$$(3.17) \quad s_k = \frac{x_k^2}{2}, \quad k = 1, \dots, d.$$

The flat metric on $\mathbb{R}^d$ becomes

$$(3.18) \quad \frac{1}{2} \sum_{k=1}^d \frac{(ds_k)^2}{s_k}.$$

Consider the moment map ν restricted to Z'_r . The above arguments show that ν maps Z'_r diffeomorphically onto the interior Σ° of Σ . In particular we have

$$(3.19) \quad \langle \nu(x), u_k \rangle = \lambda_k + s_k, \quad k = 1, \dots, d, \quad \text{for } x \in Z'_r.$$

Let $l_k : \mathbb{R}^{n^*} \rightarrow \mathbb{R}$ be the affine function

$$l_k(x) = \langle x, u_k \rangle - \lambda_k, \quad k = 1, \dots, d.$$

Then by equation (3.19) we have

$$(3.20) \quad l_k \circ \nu = s_k.$$

Thus the moment map ν pulls back the metric

$$(3.21) \quad \frac{1}{2} \sum_{k=1}^d \frac{(dl_k)^2}{l_k},$$

on Σ° to the metric (3.18) on Z'_r . We obtain the following.

Proposition 3.20. *The moment map $\nu : X'_r \rightarrow \Sigma^\circ$ is an isometry when Σ° is given the metric (3.21).*

Let $W \subset X$ be the orbit of $T_{\mathbb{C}}^n$ isomorphic to $T_{\mathbb{C}}^n$. Then by restriction W has a T^n -invariant Kähler form ω . Identify $T_{\mathbb{C}}^n = \mathbb{C}^n/2\pi i\mathbb{Z}^n$, so there is an inclusion $\iota : \mathbb{R}^n \rightarrow T_{\mathbb{C}}^n$.

Proposition 3.21. *Let ω be a T^n -invariant Kähler form on W . Then the action of T^n is Hamiltonian if and only if ω has a T^n -invariant potential function, that is, a function $F \in C^\infty(\mathbb{R}^n)$ such that*

$$\omega = 2i\partial\bar{\partial}F.$$

Proof. Suppose the action is Hamiltonian. Any T^n -orbit is Lagrangian, so ω restricts to zero. The inclusion $T^n \subset T_{\mathbb{C}}^n$ is a homotopy equivalence. Thus ω is exact. Let γ be a T^n -invariant 1-form with $\omega = d\gamma$. Let $\gamma = \beta + \bar{\beta}$ where $\beta \in \Omega^{0,1}$. Then

$$\omega = d\gamma = \partial\beta + \bar{\partial}\bar{\beta},$$

since $\bar{\partial}\beta = \partial\bar{\beta} = 0$. Since $H^{0,k}(W)_{T^n} = 0$ for $k > 0$, there exists a T^n -invariant function f with $\beta = \bar{\partial}f$. Then

$$\omega = \partial\bar{\partial}f + \bar{\partial}\partial\bar{f} = 2i\partial\bar{\partial}\text{Im } f.$$

The converse is a standard result. $\square$

Suppose the T^n action on W is Hamiltonian with moment map $\nu : W \rightarrow \mathbb{R}^{n*}$. Denote by $x + iy$ the coordinates given by the identification $W = \mathbb{C}^n/2\pi i\mathbb{Z}^n$.

Proposition 3.22 ([22]). *Up to a constant ν is the Legendre transform of F , i.e.*

$$\nu(x + iy) = \frac{\partial F}{\partial x} + c, \quad c \in \mathbb{R}^{n*}$$

Proof. By definition

$$d\nu_k = -\iota\left(\frac{\partial}{\partial y_k}\right)\omega.$$

But by Proposition 3.21,

$$\omega = \sum_{j,k=1}^n \frac{\partial^2 F}{\partial x_j \partial x_k} dx_j \wedge dy_k,$$

so

$$d\nu_k = -\iota\left(\frac{\partial}{\partial y_k}\right)\omega = d\left(\frac{\partial F}{\partial x_k}\right).$$

Therefore $\nu_k = \frac{\partial F}{\partial x_k} + c_k$. $\square$

We can eliminate c by replacing F with $F - \sum_{k=1}^n c_k x_k$.

Notice that the metric (3.21) on Σ° can be written

$$(3.22) \quad \sum_{j,k} \frac{\partial^2 G}{\partial y_j \partial y_k} dy_j dy_k,$$

with

$$(3.23) \quad G = \frac{1}{2} \sum_{k=1}^d l_k(y) \log l_k(y).$$

V. Guillemin [22] showed that the Legendre transform of G is the inverse Legendre transform of F , i.e.

$$(3.24) \quad \frac{\partial F}{\partial x} = y \text{ and } \frac{\partial G}{\partial y} = x.$$

From this it follows that

$$(3.25) \quad F(x) = \sum_{i=1}^n x_i y_i - G(y), \text{ where } y = \frac{\partial F}{\partial x}.$$

Define

$$l_\infty(x) = \sum_{i=1}^d \langle x, u_i \rangle.$$

From equations (3.23) and (3.25) it follows that F has the expression

$$(3.26) \quad F = \frac{1}{2} \nu^* \left(\sum_{k=1}^d \lambda_k \log l_k + l_\infty \right),$$

which gives us the following.

Theorem 3.23 ([22, 11]). *On the open $T_{\mathbb{C}}^n$ orbit of X_{Σ^*} the Kähler form ω is given by*

$$i\partial\bar{\partial}\nu^* \left(\sum_{k=1}^d \lambda_k \log l_k + l_\infty \right).$$

Suppose we have an embedding as in Proposition 3.16,

$$\psi_h : X_{\Sigma^*} \rightarrow \mathbb{C}P^N.$$

So Σ_h is an integral polytope and $M \cap \Sigma_h = \{m_0, m_1, \dots, m_N\}$. Let ω_{FS} be the Fubini-Study metric on $\mathbb{C}P^N$. Note that $\psi_h^* \omega_{FS}$ is degenerate along the singular set of X , so does not define a Kähler form.

Consider the restriction of ψ_h to the open $T_{\mathbb{C}}^n$ orbit $W \subset X$. Let $\iota = \psi_h|_W$. It is induced by a representation

$$(3.27) \quad \tau : T_{\mathbb{C}}^n \rightarrow GL(N+1, \mathbb{C}),$$

with weights $m_0, m_1, \dots, m_N$. If $z = x + iy \in \mathbb{C}^n / 2\pi i \mathbb{Z}^n = T_{\mathbb{C}}^n$, and $w = (w_0, \dots, w_N)$, then

$$(3.28) \quad \tau(\exp z)w = (e^{\langle m_0, x+iy \rangle} w_0, \dots, e^{\langle m_N, x+iy \rangle} w_N).$$

Recall the Fubini-Study metric is

$$(3.29) \quad \omega_{FS} = i\partial\bar{\partial} \log |w|^2.$$

Let $[w_0 : \dots : w_N]$ be homogeneous coordinates of a point in the image of W , then

$$(3.30) \quad \iota^* \omega_{FS} = i\partial\bar{\partial} \log \left(\sum_{k=0}^N |w_k|^2 e^{2\langle m_k, x \rangle} \right).$$

From equation (3.23) we have

$$x = \frac{\partial G}{\partial y} = \frac{1}{2} \left(\sum_{j=1}^d u_j \log l_j + u \right),$$

where $u = \sum u_j$. Then

$$2\langle m_i, x \rangle = 2\langle m_i, \frac{\partial G}{\partial y} \rangle = \sum_{j=1}^d \langle m_i, u_j \rangle \log l_j + \langle m_i, u \rangle.$$

So setting $d_i = e^{\langle m_i, u \rangle}$, gives

$$e^{2\langle m_i, x \rangle} = \nu^* \left(d_i \prod_{j=1}^d l_j^{\langle m_i, u_j \rangle} \right).$$

But from (3.26),

$$e^{2F} = \nu^* \left(e^{l_\infty} \prod_{j=1}^d l_j^{\lambda_j} \right).$$

Combining these,

$$e^{2\langle m_i, x \rangle} = e^{2F} \nu^* \left(d_i e^{-l_\infty} \prod_{j=1}^d l_j^{l_j(m_i)} \right).$$

Let $k_i = |w_i|^2 d_i$, then summing gives

$$\sum_{i=1}^N |w_i|^2 e^{2\langle m_i, x \rangle} = e^{2F} \nu^* (e^{-l_\infty} Q),$$

where

$$Q = \sum_{i=1}^N k_i \prod_{j=1}^d l_j^{l_j(m_i)}.$$

Thus we have

$$(3.31) \quad \psi_h^* \omega_{FS} = \omega + i\partial\bar{\partial} \nu^* (-l_\infty + \log Q).$$

Using that Σ_h is integral, and $k_i \neq 0$ for m_i a vertex of Σ_h , it is not difficult to show that Q is a positive function on Σ_h . Thus equation (3.31) is valid on all of X .

Theorem 3.24. *Suppose $\mathbf{L}_h$ is very ample for some $h \in \text{SF}(\Delta^*)$ strictly upper convex and integral, and let ω be the canonical Kähler metric for the polytope Σ_h . Then*

$$[\omega] = 2\pi c_1(\mathbf{L}) = [\psi_h^* \omega_{FS}].$$

Corollary 3.25. *Suppose $X = X_{\Delta^*}$ is Fano. Let ω be the canonical metric of the integral polytope Σ_{-k}^* . Then*

$$[\omega] = 2\pi c_1(\mathbf{K}^{-1}) = 2\pi c_1(X).$$

Thus $c_1(X) > 0$. Conversely, if $c_1(X) > 0$, then $\mathbf{K}^{-p}$ is very ample for some $p > 0$ and X is Fano as defined in definition (3.15).

Proof. For some $p \in \mathbb{Z}^+$, $-pk \in \text{SF}(\Delta^*)$ is integral and $\mathbf{L}_{-pk} = \mathbf{K}^{-p}$ is very ample. Let $\tilde{\omega}$ be the canonical metric of the integral polytope Σ_{-pk}^* . From the theorem we have

$$[\tilde{\omega}] = 2\pi c_1(\mathbf{K}^{-p}) = 2\pi p c_1(X).$$

Let ω be the canonical metric for Σ_{-k}^* . Theorem (3.23) implies that $[\tilde{\omega}] = p[\omega]$

For the converse, It follows from the extension to orbifolds of the Kodaira embedding theorem of W. Baily [2] that $\mathbf{K}^{-p}$ is very ample for some $p > 0$ sufficiently

large. It follows from standard results on toric varieties that $-k$ is strictly upper convex (see [34]). $\square$

The next result will have interesting applications to the Einstein manifolds constructed later.

Proposition 3.26. *With the canonical metric the volume of X_{Σ^*} is $(2\pi)^n$ times the Euclidean volume of Σ .*

Proof. Let $W \subset X$ be the open $T_{\mathbb{C}}^n$ orbit. We identify W with $\mathbb{C}^n/2\pi i\mathbb{Z}^n$ with coordinates $x + iy$. The restriction of ω to W is

$$\omega|_W = \sum_{j,k=1}^n \frac{\partial^2 F}{\partial x_j \partial x_k} dx_j \wedge dy_k.$$

Thus

$$\frac{\omega^n}{n!} = \det\left(\frac{\partial^2 F}{\partial x_j \partial x_k}\right) dx \wedge dy.$$

Integrating over dy gives

$$\text{Vol}(X, \omega) = (2\pi)^n \int_{\mathbb{R}^n} \det\left(\frac{\partial^2 F}{\partial x_j \partial x_k}\right) dx.$$

By Proposition 3.22 $x \rightarrow z = \nu(x + iy) = \frac{\partial F}{\partial x}$ is a diffeomorphism from $\mathbb{R}^n$ to Σ° . By the change of variables,

$$\text{Vol}(\Sigma) = \int_{\Sigma} dz = \int_{\mathbb{R}^n} \det\left(\frac{\partial^2 F}{\partial x_j \partial x_k}\right) dx.$$

$\square$

Corollary 3.27. *Let $X = X_{\Delta^*}$ be a toric Fano orbifold. And let ω be any Kähler form with $\omega \in c_1(X)$. Then*

$$\text{Vol}(X, \omega) = \frac{1}{n!} c_1(X)^n [X] = \text{Vol}(\Sigma_{-k}).$$

Proof. Let ω_c be the canonical metric associated to Σ_{-k}^* , then $\frac{1}{2\pi}\omega_c \in c_1(X)$ by Corollary 3.25. Then

$$\text{Vol}(X, \omega) = \frac{1}{(2\pi)^n} \text{Vol}(X, \omega_c) = \text{Vol}(\Sigma_{-k}).$$

$\square$

3.3. Moment map and Futaki invariant. A closer analysis of the moment map in the Fano case, originally due to T. Mabuchi [30], will be useful. Suppose in this section that $X = X_{\Delta^*}$ is toric Fano with $\dim_{\mathbb{C}} X = n$. As above, $z_k = x_k + iy_k, 1 \leq k \leq n$ are logarithmic coordinates, $(z_1, \dots, z_n) \in \mathbb{C}^n/2\pi i\mathbb{Z}^n \cong T_{\mathbb{C}}$. For $(t_1, \dots, t_n) \in T_{\mathbb{C}}$, $x_k = \log |t_k|$. Then a T -invariant function $u \in C^\infty(T_{\mathbb{C}})$ is considered as a C^∞ function $u = u(x_1, \dots, x_n)$ on $\mathbb{R}^n$. There exists a T -invariant fiber metric Ω on $\mathbf{K}_X^{-1}$ with positive Chern form. Thus there exists a C^∞ function $u = u(x_1, \dots, x_n)$ so that

$$(3.32) \quad e^{-u} \prod_{k=1}^n (dx_k \wedge dy_k),$$

extends to a volume form Ω on all of X and $i\partial\bar{\partial}u$ extends to a Kähler form ω . The moment map $\nu_u : X \rightarrow M_{\mathbb{R}}$ can be given, without an ambiguous constant, as

$$(3.33) \quad \nu_u(t) = \left(\frac{\partial u}{\partial x_1}(t), \dots, \frac{\partial u}{\partial x_n}(t) \right), \quad \text{for } t \in T_{\mathbb{C}}.$$

Theorem 3.28. *The closure of the image $\nu_u(T_{\mathbb{C}})$ in $M_{\mathbb{R}}$ is Σ_{-k} . Furthermore, ν_u extends to a C^∞ map $\nu_u : X \rightarrow M_{\mathbb{R}}$, which is the usual moment map.*

Only the first statement remains to be proved. This is a slight generalization of a similar result in [30].

We define the Futaki invariant. Let $\text{Aut}^o(X) \subseteq \text{Aut}(X)$ be the subgroup of the homomorphic automorphism group preserving the orbifold structure, and let $\mathfrak{g}$ be its Lie algebra. Let $\omega \in 2\pi c_1(X)$ be a Kähler form. There exists $f \in C^\infty(X)$ with $\text{Ricci}(\omega) - \omega = i\partial\bar{\partial}f$. Set $c = -2^{n+1}((2\pi c_1(X))^n[X])^{-1}$. Then the *Futaki invariant* $F : \mathfrak{g} \rightarrow \mathbb{C}$ is defined by

$$(3.34) \quad F(V) = c \int_X V f \omega^n, \quad \text{for } V \in \mathfrak{g}.$$

Note that, as proved in [17], F is zero on $[\mathfrak{g}, \mathfrak{g}]$. We have the Cartan decomposition

$$\mathfrak{g} = \mathfrak{t}_{\mathbb{C}} \oplus \sum_i \mathbb{C}v_i,$$

where the v_i are eigenvalues for the adjoint action of $\mathfrak{t}_{\mathbb{C}}$. Since $\mathfrak{t}$ is the Lie algebra of a maximal torus, one see that the v_i are contained in $[\mathfrak{g}, \mathfrak{g}]$. Thus we may restrict F to $\mathfrak{t}_{\mathbb{C}}$.

Suppose $\mathbf{L}$ is an equivariant holomorphic line V-bundle on X . In our case of interest $\mathbf{L} = \mathbf{K}_X^{-1}$ with the usual action. Let H be the space of hermitian metrics on $\mathbf{L}$. For $h \in H$ denote by $c_1(\mathbf{L}, h)$ the Chern form, $\frac{i}{2\pi}\bar{\partial}\partial \log(h)$ in local holomorphic coordinates. For a pair $(h', h'') \in H \times H$ we define

$$R_{\mathbf{L}}(h', h'') := \int_{t_0}^{t_1} \left(\int_X h_t^{-1} \dot{h}_t (2\pi c_1(\mathbf{L}, h_t))^n \right) dt,$$

where $h_t, t_0 \leq t_1$ is any piecewise smooth path with $h_{t_0} = h'$ and $h_{t_1} = h''$. One has that $R_{\mathbf{L}}(h', h'')$ is independent of the path h_t and satisfies

$$R_{\mathbf{L}}(g^*h', g^*h'') = R_{\mathbf{L}}(h', h''), \quad \text{for } g \in \text{Aut}^o(X),$$

and the cocycle conditions

$$(3.35) \quad R_{\mathbf{L}}(h', h'') + R_{\mathbf{L}}(h'', h') = 0, \text{ and}$$

$$(3.36) \quad R_{\mathbf{L}}(h, h') + R_{\mathbf{L}}(h', h'') + R_{\mathbf{L}}(h'', h) = 0,$$

for any $h, h', h'' \in H$. These identities imply that

$$(3.37) \quad r_{\mathbf{L}}(g) := \exp(R_{\mathbf{L}}(h, g^*h)), \quad \text{for } g \in \text{Aut}^o(X),$$

is independent of $h \in H$ and is a Lie group homomorphism into $\mathbb{R}_+$. It has associated Lie algebra character $r_{\mathbf{L}*} : \mathfrak{g} \rightarrow \mathbb{R}$.

Let σ be a $T_{\mathbb{C}}$ -invariant section of $\mathbf{L}^*$. Then $h \in H$ is $h = e^{-u_h} \sigma \otimes \bar{\sigma}$ on $W \cong T_{\mathbb{C}}$ for some $u_h \in C^\infty(W)$. We denote by $V_{\mathbb{R}}$ the real component of a homomorphic vector field. Then differentiating (3.37) gives

$$(3.38) \quad r_{\mathbf{L}*}(V) = - \int_W V_{\mathbb{R}}(u_h) (i\partial\bar{\partial}u_h)^n,$$

for $V \in \mathfrak{t}_{\mathbb{C}}$ independent of $h \in H$. We have the following.

Proposition 3.29. *Suppose X_{Δ^*} is a toric Fano orbifold. Then with $\mathbf{L} = \mathbf{K}_X^{-1}$ we have*

$$F_X = -2^{n+1}((2\pi c_1(X))^n[X])^{-1}r_{\mathbf{L}*},$$

where both sides are restricted to $\mathfrak{t}_{\mathbb{C}}$.

Proof. By assumption there is an $h \in H$ with positive Chern form. Let β be a $T_{\mathbb{C}}$ -invariant section of $\mathbf{K}_X^{-1}$ over W . Then h may be written as the volume form on W $\Omega = i^n(-1)^{n(n-1)/2}e^{-u_h}\beta \wedge \bar{\beta}$ and $i\partial\bar{\partial}u_h$ extends to a Kähler form $\omega \in 2\pi c_1(X)$. Then if $f = \log(\frac{\Omega}{\omega^n})$, we have

$$\text{Ricci}(\omega) - \omega = i\partial\bar{\partial}f.$$

Then

$$\begin{aligned} 0 &= \int_X \mathcal{L}_{V_{\mathbb{R}}}(e^{-f}\Omega) \\ &= - \int_X V_{\mathbb{R}}(f)\omega^n + \int_X e^{-f} \mathcal{L}_{V_{\mathbb{R}}}\Omega \\ &= - \int_X V_{\mathbb{R}}(f)\omega^n - \int_W V_{\mathbb{R}}(u_h)\omega^n \end{aligned}$$

And the result follows from (3.38). $\square$

The $t_k \frac{\partial}{\partial t_k}, k = 1, \dots, n$ form a basis of $\mathfrak{t}_{\mathbb{C}}$. We may assume that the Kähler form ω is T -invariant. Then we have

$$\begin{aligned} F_X(t_k \frac{\partial}{\partial t_k}) &= -2^{n+1}((2\pi c_1(X))^n[X])^{-1}r_{\mathbf{L}*}(t_k \frac{\partial}{\partial t_k}) \\ &= 2^n((2\pi c_1(X))^n[X])^{-1} \int_W \frac{\partial u_h}{\partial x_k} (i\partial\bar{\partial}u_h)^n \\ &= 2^n((2\pi c_1(X))^n[X])^{-1} \int_W \frac{\partial u_h}{\partial x_k} \left(\frac{1}{2}\right)^n n! \det\left(\frac{\partial^2 u_h}{\partial x_j \partial x_k}\right) \prod_{l=1}^n dx_l \wedge dy_l \\ &= (2\pi)^n((2\pi c_1(X))^n[X])^{-1} \int_{\mathbb{R}^n} \frac{\partial u_h}{\partial x_k} n! \det\left(\frac{\partial^2 u_h}{\partial x_j \partial x_k}\right) dx \\ &= (2\pi)^n n!((2\pi c_1(X))^n[X])^{-1} \int_{\Sigma_{-k}} y_k dy \\ &= \text{Vol}(\Sigma_{-k})^{-1} \int_{\Sigma_{-k}} y_k dy \end{aligned}$$

where $y_k = \frac{\partial u_h}{\partial x_k}$.

We have the following simple interpretation of the Futaki invariant in this case.

Proposition 3.30. *Suppose $X = X_{\Delta^*}$ is a toric Fano orbifold. Then the Futaki invariant F_X is the barycenter of the polytope Σ_{-k} .*

3.4. Symmetric toric orbifolds. Let X_{Δ} be an n -dimensional toric variety. Let $\mathcal{N}(T_{\mathbb{C}}) \subset \text{Aut}(X)$ be the normalizer of $T_{\mathbb{C}}$. Then $\mathcal{W}(X) := \mathcal{N}(T_{\mathbb{C}})/T_{\mathbb{C}}$ is isomorphic to the finite group of all symmetries of Δ , i.e. the subgroup of $GL(n, \mathbb{Z})$ of all $\gamma \in GL(n, \mathbb{Z})$ with $\gamma(\Delta) = \Delta$. Then we have the exact sequence.

$$(3.39) \quad 1 \rightarrow T_{\mathbb{C}} \rightarrow \mathcal{N}(T_{\mathbb{C}}) \rightarrow \mathcal{W}(X) \rightarrow 1.$$

Choosing a point $x \in X$ in the open orbit, defines an inclusion $T_{\mathbb{C}} \subset X$. This also provides a splitting of (3.39). Let $\mathcal{W}_0(X) \subseteq \mathcal{W}(X)$ be the subgroup which are also automorphisms of Δ^* ; $\gamma \in \mathcal{W}_0(X)$ is an element of $\mathcal{N}(T_{\mathbb{C}}) \subset \text{Aut}(X)$ which preserves the orbifold structure. Let $G \subset \mathcal{N}(T_{\mathbb{C}})$ be the compact subgroup generated by T^n , the maximal compact subgroup of $T_{\mathbb{C}}$, and $\mathcal{W}_0(X)$. Then we have the, split, exact sequence

$$(3.40) \quad 1 \rightarrow T^n \rightarrow G \rightarrow \mathcal{W}_0(X) \rightarrow 1.$$

Definition 3.31. *A symmetric Fano toric orbifold X is a Fano toric orbifold with $\mathcal{W}_0$ acting on N with the origin as the only fixed point. Such a variety and its orbifold structure is characterized by the convex polytope Δ^* invariant under $\mathcal{W}_0$. We call a toric orbifold special symmetric if $\mathcal{W}_0(X)$ contains the involution $\sigma : N \rightarrow N$, where $\sigma(n) = -n$.*

The following is immediate from Proposition 3.30.

Proposition 3.32. *For a symmetric Fano toric orbifold X one has $F_X \equiv 0$.*

Definition 3.33. *The index of a Fano orbifold X is the largest positive integer m such that there is a holomorphic V -bundle $\mathbf{L}$ with $\mathbf{L}^m \cong \mathbf{K}_X^{-1}$. The index of X is denoted $\text{Ind}(X)$.*

Note that $c_1(X) \in H_{orb}^2(X, \mathbb{Z})$, and $\text{Ind}(X)$ is the greatest positive integer m such that $\frac{1}{m}c_1(X) \in H_{orb}^2(X, \mathbb{Z})$.

Proposition 3.34. *Let X_{Δ^*} be a special symmetric toric Fano orbifold. Then $\text{Ind}(X) = 1$ or 2 .*

Proof. We have $\mathbf{K}^{-1} \cong \mathbf{L}_{-k}$ with $-k \in \text{SF}(\Delta^*)$ where $-k(n_\rho) = -1$ for all $\rho \in \Delta(1)$. Suppose we have $\mathbf{L}^m \cong \mathbf{K}^{-1}$. By Proposition 3.12 there is an $h \in \text{SF}(\Delta^*)$ and $f \in M$ so that $mh = -k + f$. For some $\rho \in \Delta(1)$,

$$\begin{aligned} mh(n_\rho) &= -1 + f(n_\rho) \\ mh(-n_\rho) &= -1 - f(n_\rho). \end{aligned}$$

Thus $m(h(n_\rho) + h(-n_\rho)) = -2$, and $m = 1$ or 2 . $\square$

We will now restrict to dimension two. In the smooth case every Fano surface, called a *del Pezzo surface*, is either $\mathbb{CP}^1 \times \mathbb{CP}^1$ or $\mathbb{CP}^2$ blown up at r points in general position $0 \leq r \leq 8$. The smooth toric Fano surfaces are $\mathbb{CP}^1 \times \mathbb{CP}^1$, $\mathbb{CP}^2$, the Hirzebruch surface F_1 , the equivariant blow up of $\mathbb{CP}^2$ at two $T_{\mathbb{C}}$ -fixed points, and the equivariant blow up of $\mathbb{CP}^2$ at three $T_{\mathbb{C}}$ -fixed points. There are only three examples of smooth symmetric toric Fano surfaces, which are $\mathbb{CP}^1 \times \mathbb{CP}^1$, $\mathbb{CP}^2$, and the equivariant blow up of $\mathbb{CP}^2$ at three $T_{\mathbb{C}}$ -fixed points. Their marked fans are shown in figure 1. The smooth toric Fano surfaces admitting a Kähler-Einstein metric are precisely the symmetric cases.

4. EINSTEIN EQUATION

We consider the existence of Sasaki-Einstein metrics on toric Sasakian manifolds. This problem is completely solved in [18] where the more generally the existence of Sasaki-Ricci solitons is proved extending the existence of Kähler-Ricci solitons on toric Fano manifolds proved in [41]. For the examples of Sasaki-Einstein manifolds considered in this article a complete proof can be found in [40].




FIGURE 1. The three smooth examples




FIGURE 2. Example with 8 point singular set and $\mathcal{W}_0 = \mathbb{Z}_2$

Given a Sasakian manifold (M, g, Φ, ξ, η) we consider deformations of the transversal Kähler structure. That is, for a basic function $\phi \in C_B^\infty(M)$, set

$$\tilde{\eta} = \eta + 2d_B^c \phi.$$

Then for ϕ small enough,

$$\tilde{\omega}^T = \frac{1}{2}d\tilde{\eta} = \frac{1}{2}d\eta + d_B d_B^c \phi = \omega^T + d_B d_B^c \phi$$

is a transversal Kähler metric and $\tilde{\eta} \wedge \tilde{\eta}^m$ is nowhere zero. Then there is a Sasakian structure $(M, \tilde{g}, \tilde{\Phi}, \xi, \tilde{\eta})$ with the same Reeb vector field, transverse holomorphic structure, and basic Kähler class $[\tilde{\omega}^T]_B = [\omega^T]_B$. The existence of such a deformation of the transverse Kähler structure to a Sasakian Einstein structure requires the following.

Proposition 4.1. *The following three conditions are equivalent.*

- (i) $(2m+2)\omega \in 2\pi c_1^B(M)$.
- (ii) $\mathbf{K}_{C(M)}^q$ is holomorphically trivial for some $q \in \mathbb{Z}_+$, and there is a nowhere zero section Ω of $\mathbf{K}_{C(M)}$, which will be multi-valued if $\mathbf{K}_{C(M)}^q$ is not trivial, with $\mathcal{L}_{r\partial_r}\Omega = (m+1)\Omega$.
- (iii) $\mathbf{K}_{C(M)}^q$ is holomorphically trivial for some $q \in \mathbb{Z}_+$, and there is a section Ω of $\mathbf{K}_{C(M)}$, which will be multi-valued if $\mathbf{K}_{C(M)}^q$ is not trivial, such that

$$\frac{i^{m+1}}{2^{m+1}}(-1)^{\frac{m(m+1)}{2}}\Omega \wedge \bar{\Omega} = e^f \frac{1}{n!}\bar{\omega}^{m+1},$$

for $f \in C_B^\infty(M)$ pulled back to an element of $C^\infty(C(M))$. Here $\bar{\omega}$ is the Kähler form on $C(M)$.

In Proposition 2.4 with $\mathbf{L} = \mathbf{K}_X^{\frac{1}{2}}$ the conditions of Proposition 4.1 are satisfied. We will need the following definition.

Definition 4.2. A Hamiltonian holomorphic vector field on M is a complex vector field Y invariant by ξ so that

- (i) For any local leaf space projection $\pi_\alpha : U_\alpha \rightarrow V_\alpha$, $\pi_\alpha(Y)$ is a holomorphic vector field,
- (ii) the complex function $\theta_Y = \sqrt{-1}\eta(Y)$ satisfies

$$\bar{\partial}_B\theta_Y = -\frac{\sqrt{-1}}{2}Y\lrcorner d\eta.$$

The Lie algebra of Hamiltonian holomorphic vector fields on M is denoted $\mathfrak{h}$.

Note that $\tilde{Y} = Y + i\eta(Y)r\partial_r$ is a holomorphic vector field on $C(M)$. These correspond exactly to transversely holomorphic vector fields, i.e. those satisfying (i), which have a potential function (cf. [10]). In other words if $Y \in \Gamma(D \otimes \mathbb{C})$ satisfies (i) and $\mathcal{L}_\xi Y = 0$, and there exists a complex function $\theta_Y \in C_B^\infty(M)$ with

$$\bar{\partial}_B\theta_Y = -\frac{\sqrt{-1}}{2}Y\lrcorner d\eta,$$

then $Y - \sqrt{-1}\theta_Y\xi$ is a Hamiltonian holomorphic vector field. Furthermore, in the case $c_1^B(M) > 0$ by the transverse Calabi-Yau theorem (cf. [28]) there exists a transversal Kähler deformation $(M, \tilde{g}, \tilde{\Phi}, \xi, \tilde{\eta})$ with $\tilde{\eta} = \eta + 2d_B^c\phi$ for some basic $\phi \in C_B^\infty(M)$ with $\text{Ric}_{\tilde{g}^T}$ positive. The usual Weitzenböck formula shows that the space of basic harmonic 1-forms $\mathcal{H}_B^1$ is zero. It follows that if $Y \in \Gamma(D \otimes \mathbb{C})$ satisfies (i) and $\mathcal{L}_\xi Y = 0$, there exists a potential function $\theta_Y \in C_B^\infty(M)$ so that $Y - \sqrt{-1}\theta_Y\xi$ is a Hamiltonian holomorphic vector field. Thus $\mathfrak{h}$ is isomorphic to the space of transversely holomorphic vector fields commuting with ξ , which is isomorphic to the space of holomorphic vector fields on $C(M)$ commuting with $\xi + \sqrt{-1}r\partial_r$.

We now suppose that the conditions of Proposition 4.1 hold for (M, g, Φ, ξ, η) . Let Y be a Hamiltonian holomorphic vector field. Then (M, g, Φ, ξ, η) is a *Sasakian-Ricci soliton* if

$$(4.1) \quad \text{Ric}^T - (2m+2)g^T = \mathcal{L}_Y g^T.$$

Let $h \in C_B^\infty(M)$ be a basic function with

$$(4.2) \quad \text{Ricci}(\omega^T) - (2m+2)\omega^T = i\partial_B\bar{\partial}_B h,$$

where $\omega^T = \frac{1}{2}d\eta$ is the transverse Kähler form. In [39] Tian and Zhu defined a modified Futaki invariant F_Y

$$(4.3) \quad F_Y(v) = \int_M v(h - \theta_Y) e^{\theta_Y} \eta \wedge \left(\frac{1}{2}d\eta\right)^m, \quad v \in \mathfrak{h}.$$

One can show as in [39] that F_Y is unchanged under transversal deformations $\eta \rightarrow \tilde{\eta} + 2d_B^c \phi$ of the Sasakian structure; and $F_Y(v) = 0$, for all $v \in \mathfrak{h}$, is a necessary condition for a solution to (4.1). If θ_Y is a constant, i.e. $Y = c\xi$ for some $c \in \mathbb{C}$, then (4.3) defines the usual Futaki invariant. And if M is quasi-regular this is, up to a constant, the same invariant defined in (3.34).

Let H be the transversal holomorphic automorphism group generated by $\mathfrak{h}$. Let $K \subset H$ be a compact group with Lie algebra $\mathfrak{k}$, and let $\mathfrak{k}^\mathbb{C}$ be its complexification. Note that we may choose a K -invariant Sasakian structure $(M, \tilde{g}, \tilde{\Phi}, \xi, \tilde{\eta})$. Then for $Y \in \mathfrak{k}$, one can take θ_Y to be imaginary. So one has $\mathfrak{k}^\mathbb{C} \subset \mathfrak{h}$. As in [39] we have the following.

Proposition 4.3. *There exists a $Y \in \mathfrak{k}^\mathbb{C}$ with $\text{Im } Y \in \mathfrak{k}$ so that*

$$F_Y(v) = 0, \quad \text{for all } v \in \mathfrak{k}^\mathbb{C}.$$

Furthermore, Y is unique up to addition of $c\xi$, for $c \in \mathbb{C}$, and

$$F_Y([v, w]) = 0, \quad \text{for all } v \in \mathfrak{k}^\mathbb{C} \text{ and } w \in \mathfrak{h}.$$

Suppose now that (M, g, Φ, ξ, η) is a toric Sasakian manifold. If $\mathfrak{t}$ is the Lie algebra of the $m+1$ -torus T acting on M , then $\mathfrak{t} \subset \mathfrak{t}_\mathbb{C} \subseteq \mathfrak{h}$. Using the same argument as after (3.34) we have the following.

Corollary 4.4. *If M is toric, then there exists a unique $Y \in \mathfrak{t}_\mathbb{C}$ with $\text{Im } Y \in \mathfrak{t}$ so that*

$$F_Y(v) = 0, \quad \text{for all } v \in \mathfrak{h}.$$

For $\phi \in C_B^\infty(M)$, $\tilde{\eta} = \eta + 2d_B^c \phi$ defines a transversally deformed Sasakian structure, with transverse Kähler form $\tilde{\omega}^T = \omega + i\partial\bar{\partial}\phi$. And the Hamiltonian function for $Y \in \mathfrak{h}$ becomes $\tilde{\theta}_Y = \theta_Y + Y\phi$ (see [39]). In transverse holomorphic coordinates (4.1) becomes the Monge-Ampère equation

$$(4.4) \quad \frac{\det(g_{i\bar{j}}^T + \phi_{i\bar{j}})}{\det(g_{i\bar{j}}^T)} = e^{-(2m+2)\phi - \theta_Y - Y\phi + h}.$$

In the toric case A. Futaki, H. Ono, and G. Wang [18] prove the necessary C^0 estimate on ϕ to solve (4.4) using the continuity method.

Theorem 4.5. *Let (M, g, Φ, ξ, η) be a compact toric Sasakian manifold satisfying the conditions in (4.1). Then there exists a unique transversal deformation $(M, \tilde{g}, \tilde{\Phi}, \xi, \tilde{\eta})$ which is a Sasakian-Ricci soliton.*

If the Futaki invariant vanishes, then one has $Y = 0$ in Corollary 4.4. Therefore we have the following.


Corollary 4.6. *The solution in Theorem 4.5 is Einstein if, and only if, the Futaki invariant vanishes.*

If (M, g, Φ, ξ, η) is quasi-regular then the leaf space of $\mathcal{F}_\xi$ is a toric orbifold $X = X_{\Sigma_{-k}^*}$ where Σ_{-k}^* is the marked polytope of the anti-canonical bundle. Then (M, g, Φ, ξ, η) admits a transversal deformation to an Sasaki-Einstein structure if,

and only if, the barycenter of Σ_{-k}^* is the origin. In particular, if M is quasi-regular and the leaf space X_{Δ^*} is symmetric, then solution is Einstein.

5. 3-SASAKIAN MANIFOLDS

In this section we define 3-Sasakian manifolds, the closely related quaternionic-Kähler spaces, and their twistor spaces. For more details see [5]. These are sister geometries where one is able to pass from one to the other two by considering the appropriate orbifold fibration. Given a 3-Sasakian manifold $\mathcal{S}$ there is the associated twistor space $\mathcal{Z}$, quaternionic-Kähler orbifold $\mathcal{M}$, and hyperkähler cone $C(\mathcal{S})$. This is characterized by the *diamond*:



The equivalent 3-Sasakian and quaternionic-Kähler reduction procedures provide an elementary method for constructing 3-Sasakian and quaternionic-Kähler orbifolds (cf. [19, 8]). This method is effective in producing smooth 3-Sasakian manifolds, though the quaternionic-Kähler spaces obtained are rarely smooth. In particular, we are interested in toric 3-Sasakian 7-manifolds $\mathcal{S}$ and their associated four dimensional quaternionic-Kähler orbifolds $\mathcal{M}$. Here toric means that the structure is preserved by an action of the real two torus T^2 . In four dimensions quaternionic-Kähler means that $\mathcal{M}$ is Einstein and anti-self-dual, i.e. the self-dual half of the Weyl curvature vanishes $W_+ \equiv 0$. These examples are well known and they are all obtained by reduction (cf. [8] and [12]). In this case we will associate two more Einstein spaces to the four Einstein spaces in the diamond. To each diamond of a toric 3-Sasakian manifold we have a special symmetric toric Fano surface X and a Sasaki-Einstein manifold M which complete diagram 1.1.

The motivation is two fold. First, it adds two more Einstein spaces to the examples on the right considered by C. Boyer, K. Galicki, and others in [8, 5] and also by D. Calderbank and M. Singer [12]. Second, M is smooth when the 3-Sasakian space $\mathcal{S}$ is. And the smoothness of $\mathcal{S}$ is ensured by a relatively mild condition on the moment map. Thus we get infinitely many quasi-regular Sasaki-Einstein 5-manifolds with arbitrarily high second Betti numbers paralleling the 3-Sasakian manifolds constructed in [8].

5.1. Definitions and basic properties. We cover some of the basics of 3-Sasakian manifolds and 3-Sasakian reduction. See [5] for more details.

Definition 5.1. Let $(\mathcal{S}, g)$ be a Riemannian manifold of dimension $n = 4m + 3$. Then $\mathcal{S}$ is 3-Sasakian if it admits three Killing vector fields $\{\xi^1, \xi^2, \xi^3\}$ each satisfying definition (2.1) such that $g(\xi^i, \xi^k) = \delta_{ij}$ and $[\xi^i, \xi^j] = 2\epsilon_{ijk}\xi^k$.

We have a triple of Sasakian structures on $\mathcal{S}$. For $i = 1, 2, 3$ we have $\eta^i(X) = g(\xi^i, X)$ and $\Phi^i(X) = \nabla_X \xi^i$. We say that $\{g, \Phi^i, \xi^i, \eta^i : i = 1, 2, 3\}$ defines a 3-Sasakian structure on $\mathcal{S}$.

Proposition 5.2. The tensors $\Phi^i, i = 1, 2, 3$ satisfy the following identities.

- (i) $\Phi^i(\xi^j) = -\epsilon_{ijk}\xi^k$,
- (ii) $\Phi^i \circ \Phi^j = -\epsilon_{ijk}\Phi^k + \xi^i \otimes \eta^j - \delta_{ij}Id$

Notice that if $\alpha = (a_1, a_2, a_3) \in S^2 \subset \mathbb{R}^3$ then $\xi(\alpha) = a_1\xi^1 + a_2\xi^2 + a_3\xi^3$ is a Sasakian structure. Thus a 3-Sasakian manifold come equipped with an S^2 of Sasakian structures.

As in definition 2.1, 3-Sasakian manifolds can be characterized by the holonomy of the cone $C(\mathcal{S})$.

Proposition 5.3. *Let $(\mathcal{S}, g)$ be a Riemannian manifold of dimension $n = 4m + 3$. Then $(\mathcal{S}, g)$ is 3-Sasakian if, and only if, the holonomy of the metric cone $(C(\mathcal{S}), \bar{g})$ is a subgroup of $Sp(m + 1)$. In other words, $(C(\mathcal{S}), \bar{g})$ is hyperkähler.*

Proof. Define almost complex structures $I_i, i = 1, 2, 3$ by

$$I_i X = -\Phi^i(X) + \eta^i(X)r\partial_r, \text{ and } I_i r\partial_r = \xi^i.$$

It is straight forward to verify that they satisfy $I_i \circ I_j = \epsilon_{ijk}I_k - \delta_{ij}Id$. And from the integrability condition on each Φ^i in definition (2.1) each $I_i, i = 1, 2, 3$ is parallel. $\square$

Since a hyperkähler manifold is Ricci flat, we have the following.

Corollary 5.4. *A 3-Sasakian manifold $(\mathcal{S}, g)$ of dimension $n = 4m + 3$ is Einstein with positive scalar curvature $s = 2(2m + 1)(4m + 3)$. Furthermore, if $(\mathcal{S}, g)$ is complete, then it is compact with finite fundamental group.*

The structure group of a 3-Sasakian manifold reduces to $Sp(m) \times \mathbb{I}_3$ where $\mathbb{I}_3$ is the 3×3 identity matrix. Thus we have

Corollary 5.5. *A 3-Sasakian manifold (M, g) is spin.*

Suppose $(\mathcal{S}, g)$ is compact. This will be the case in all examples considered here. Then the vector fields $\{\xi^1, \xi^2, \xi^3\}$ are complete and define a locally free action of $Sp(1)$ on $(\mathcal{S}, g)$. This defines a foliation $\mathcal{F}_3$, the *3-Sasakian foliation*. The generic leaf is either $SO(3)$ or $Sp(1)$, and all the leaves are compact. So $\mathcal{F}_3$ is quasi-regular, and the space of leaves is a compact orbifold, denoted $\mathcal{M}$. The projection $\varpi : \mathcal{S} \rightarrow \mathcal{M}$ exhibits $\mathcal{S}$ as an $SO(3)$ or $Sp(1)$ V-bundle over $\mathcal{M}$. The leaves of $\mathcal{F}_3$ are constant curvature 3-Sasakian 3-manifolds which must be homogeneous spherical space forms. Thus a leaf is $\Gamma \backslash S^3$ with $\Gamma \subset Sp(1)$.

For $\beta \in S^2$ we also have the characteristic vector field ξ_β with the associated 1-dimensional foliation $\mathcal{F}_\beta \subset \mathcal{F}_3$. In this case $\mathcal{F}_\beta$ is automatically quasi-regular. Denote the leaf space of $\mathcal{F}_\beta$ as $\mathcal{Z}_\beta$ or just $\mathcal{Z}$. Then the natural projection $\pi : \mathcal{S} \rightarrow \mathcal{Z}$ is an S^1 V-bundle. And $\mathcal{Z}$ is a $(2m + 1)$ -dimensional projective, normal algebraic variety with orbifold singularities and a Kähler form $\omega \in c_1^{orb}(\mathcal{Z})$, i.e. is Fano.

Fix a Sasakian structure $\{\Phi^1, \xi^1, \eta^1\}$ on $\mathcal{S}$. The horizontal subbundle $\mathcal{H} = \ker \eta^1$ to the foliation $\mathcal{F}$ of ξ^1 with the almost complex structure $I = -\Phi^1|_{\mathcal{H}}$ define a CR structure on $\mathcal{S}$. The form $\eta = \eta^2 + i\eta^3$ is of type $(1, 0)$ with respect to I . And $d\eta|_{\mathcal{H} \cap \ker(\eta)} \in \Omega^{2,0}(\mathcal{H} \cap \ker(\eta))$ is nondegenerate as a complex 2-form on $\mathcal{H} \cap \ker(\eta)$. Consider the complex 1-dimensional subspace $P \subset \Lambda^{1,0}\mathcal{H}$ spanned by η . Letting $\exp(it\xi^1)$ denote an element of the circle subgroup $U(1) \subset Sp(1)$ generated by ξ^1 one see that $\exp(it\xi^1)$ acts on P with character e^{-2it} . Then $\mathbf{L} \cong \mathcal{S} \times_{U(1)} P$ defines a holomorphic line V-bundle over $\mathcal{Z}$. And we have a holomorphic section θ of $\Lambda^{1,0}(\mathcal{Z}) \otimes \mathbf{L}$ such that

$$\theta(X) = \eta(\tilde{X}),$$

where $\tilde{X}$ is the horizontal lift of a vector field X on $\mathcal{Z}$. Let $D = \ker(\theta)$ be the complex distribution defined by θ . Then $d\theta|_D \in \Gamma(\Lambda^2 D \otimes \mathbf{L})$ is nondegenerate. Thus $D = \ker(\theta)$ is complex contact structure on $\mathcal{Z}$, that is, a maximally non-integrable holomorphic subbundle of $T^{1,0}\mathcal{Z}$. Also, $\theta \wedge (d\theta)^m$ is a nowhere zero section of $\mathbf{K}_{\mathcal{Z}} \otimes \mathbf{L}^{m+1}$. Thus $\mathbf{L} \cong \mathbf{K}_{\mathcal{Z}}^{-\frac{1}{m+1}}$ as holomorphic line V-bundles. We have the following for 3-Sasakian manifolds.

Theorem 5.6. *Let $(\mathcal{S}, g)$ be a compact 3-Sasakian manifold of dimension $n = 4m + 3$, and let $\mathcal{Z}_{\beta}$ be the leaf space of the foliation $\mathcal{F}_{\beta}$ for $\beta \in S^2$. Then $\mathcal{Z}_{\beta}$ is a compact $\mathbb{Q}$ -factorial contact Fano variety with a Kähler-Einstein metric h with scalar curvature $s = 8(2m + 1)(m + 1)$. The projection $\pi : \mathcal{S} \rightarrow \mathcal{Z}$ is an orbifold Riemannian submersion with respect to the metrics g on $\mathcal{S}$ and h on $\mathcal{Z}$.*

The space $\mathcal{Z} = \mathcal{Z}_{\beta}$ is, up to isomorphism of all structures, independent of $\beta \in S^2$. We call $\mathcal{Z}$ the *twistor space* of $\mathcal{S}$. Consider again the natural projection $\varpi : \mathcal{S} \rightarrow \mathcal{M}$ coming from the foliation $\mathcal{F}_3$. This factors into $\pi : \mathcal{S} \rightarrow \mathcal{Z}$ and $\rho : \mathcal{Z} \rightarrow \mathcal{M}$. The generic fibers of ρ is a $\mathbb{CP}^1$ and there are possible singular fibers $\Gamma \backslash \mathbb{CP}^1$ which are simply connected and for which $\Gamma \subset U(1)$ is a finite group. And restricting to a fiber $\mathbf{L}|_{\mathbb{CP}^1} = \mathcal{O}(2)$, which is a V-bundle on singular fibers. Consider $g = \exp(\frac{\pi}{2}\xi^2) \in Sp(1)$ which gives an isometry of $\mathcal{S}_{\varsigma_g} : \mathcal{S} \rightarrow \mathcal{S}$ for which $\varsigma_g(\xi^1) = -\xi^1$. And ς_g descends to an anti-holomorphic isometry $\sigma : \mathcal{Z} \rightarrow \mathcal{Z}$ preserving the fibers.

We now consider the orbifold $\mathcal{M}$ more closely. Let $(\mathcal{M}, g)$ be any $4m$ dimensional Riemannian orbifold. An *almost quaternionic* structure on $\mathcal{M}$ is a rank 3 V-subbundle $\mathcal{Q} \subset \text{End}(T\mathcal{M})$ which is locally spanned by almost complex structures $\{J_i\}_{i=1,2,3}$ satisfying the quaternionic identities $J_i^2 = -Id$ and $J_1 J_2 = -J_2 J_1 = J_3$. We say that $\mathcal{Q}$ is compatible with g if $J_i^* g = g$ for $i = 1, 2, 3$. Equivalently, each $J_i, i = 1, 2, 3$ is skew symmetric.

Definition 5.7. *A Riemannian orbifold $(\mathcal{M}, g)$ of dimension $4m, m > 1$ is quaternionic Kähler if there is an almost quaternionic structure $\mathcal{Q}$ compatible with g which is preserved by the Levi-Civita connection.*

This definition is equivalent to the holonomy of $(\mathcal{M}, g)$ being contained in $Sp(1)Sp(m)$. For orbifolds this is the holonomy on $\mathcal{M} \setminus S_{\mathcal{M}}$ where $S_{\mathcal{M}}$ is the singular locus of $\mathcal{M}$. For more on quaternionic Kähler manifolds see [37]. Notice that this definition always holds on an oriented Riemannian 4-manifold ($m = 1$). This case requires a different definition. Consider the *curvature operator* $\mathcal{R} : \Lambda^2 \rightarrow \Lambda^2$ of an oriented Riemannian 4-manifold. With respect to the decomposition $\Lambda^2 = \Lambda_+^2 \oplus \Lambda_-^2$, we have

$$(5.1) \quad \mathcal{R} = \begin{pmatrix} W_+ + \frac{s}{12} & \overset{\circ}{r} \\ \overset{\circ}{r} & W_- + \frac{s}{12} \end{pmatrix},$$

where W_+ and W_- are the selfdual and anti-self-dual pieces of the Weyl curvature and $\overset{\circ}{r} = \text{Ric} - \frac{s}{4}g$ is the trace-free Ricci curvature. An oriented 4 dimensional Riemannian orbifold $(\mathcal{M}, g)$ is quaternionic Kähler if it is Einstein and anti-self-dual, meaning that $\overset{\circ}{r} = 0$ and $W_+ = 0$.

Theorem 5.8. *Let $(\mathcal{S}, g)$ be a compact 3-Sasakian manifold of dimension $n = 4m + 3$. Then there is a natural quaternionic Kähler structure on the leaf space of $\mathcal{F}_3$, $(\mathcal{M}, \check{g})$, such that the V-bundle map $\varpi : \mathcal{S} \rightarrow \mathcal{M}$ is a Riemannian submersion. Furthermore, $(\mathcal{M}, \check{g})$ is Einstein with scalar curvature $16m(m + 2)$.*

5.2. 3-Sasakian reduction. We now summarize 3-Sasakian reduction and its application to producing infinitely many 3-Sasakian 7-manifolds. In particular, we are interested in toric 3-Sasakian 7-manifolds which have a T^2 action preserving the 3-Sasakian structure. Up to coverings they are all obtainable by taking 3-Sasakian quotients of S^{4n-1} by a torus T^k , $k = n - 2$. See [8, 5] for more details.

Let $(\mathcal{S}, g)$ be a 3-Sasakian manifold. And let $I(\mathcal{S}, g)$ be the subgroup in the isometry group $\text{Isom}(\mathcal{S}, g)$ of 3-Sasakian automorphisms.

Definition 5.9. *Let $(\mathcal{S}, g)$ be a 3-Sasakian 7-manifold. Then $(\mathcal{S}, g)$ is toric if there is a real 2-torus $T^2 \subseteq I(\mathcal{S}, g)$.*

Let $G \subseteq I(\mathcal{S}, g)$ be compact with Lie algebra $\mathfrak{g}$. One can define the 3-Sasakian moment map

$$(5.2) \quad \mu_{\mathcal{S}} : \mathcal{S} \rightarrow \mathfrak{g}^* \otimes \mathbb{R}^3$$

by

$$(5.3) \quad \langle \mu_{\mathcal{S}}^a, X \rangle = \frac{1}{2} \eta^a(\tilde{X}), \quad a = 1, 2, 3 \text{ for } X \in \mathfrak{g},$$

where $\tilde{X}$ be the vector field on $\mathcal{S}$ induced by $X \in \mathfrak{g}$.

Proposition 5.10. *Let $(\mathcal{S}, g)$ be a 3-Sasakian manifold and $G \subset I(\mathcal{S}, g)$ a connected compact subgroup. Assume that G acts freely (locally freely) on $\mu_{\mathcal{S}}^{-1}(0)$. Then $\mathcal{S} // G = \mu_{\mathcal{S}}^{-1}(0)/G$ has the structure of a 3-Sasakian manifold (orbifold). Let $\iota : \mu_{\mathcal{S}}^{-1}(0) \rightarrow \mathcal{S}$ and $\pi : \mu_{\mathcal{S}}^{-1}(0) \rightarrow \mu_{\mathcal{S}}^{-1}(0)/G$ be the corresponding embedding and submersion. Then the metric $\check{g}$ and 3-Sasakian vector fields are defined by $\pi^* \check{g} = \iota^* g$ and $\pi_* \xi^i|_{\mu_{\mathcal{S}}^{-1}(0)} = \check{\xi}^i$.*

Consider the unit sphere $S^{4n-1} \subset \mathbb{H}^n$ with the metric g obtained by restricting the flat metric on $\mathbb{H}^n$. Give S^{4n-1} the standard 3-Sasakian structure induced by the right action of $Sp(1)$. Then $I(S^{4n-1}, g) = Sp(n)$ acting by the standard linear representation on the left. We have the maximal torus $T^n \subset Sp(n)$ and every representation of a subtorus T^k is conjugate to an inclusion $\iota_{\Omega} : T^k \rightarrow T^n$ which is represented by a matrix

$$(5.4) \quad \iota_{\Omega}(\tau_1, \dots, \tau_k) = \begin{pmatrix} \prod_{i=1}^k \tau_1^{a_i^1} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \prod_{i=1}^k \tau_n^{a_i^n} \end{pmatrix},$$

where $(\tau_1, \dots, \tau_k) \in T^k$. Every such representation is defined by the $k \times n$ integral weight matrix

$$(5.5) \quad \Omega = \begin{pmatrix} a_1^1 & \cdots & a_k^1 & \cdots & a_n^1 \\ a_1^2 & \cdots & a_k^2 & \cdots & a_n^2 \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ a_1^k & \cdots & a_k^k & \cdots & a_n^k \end{pmatrix}$$

Let $\{e_i\}, i = 1, \dots, k$ be a basis for the dual of the Lie algebra of T^k , $\mathfrak{t}_k^* \cong \mathbb{R}^k$. Then the moment map $\mu_{\Omega} : S^{4n-1} \rightarrow \mathfrak{t}_k^* \otimes \mathbb{R}^3$ can be written as $\mu_{\Omega} = \sum_j \mu_{\Omega}^j e_j$, where in terms of complex coordinates $u_l = z_l + w_l j$ on $\mathbb{H}^n$

$$(5.6) \quad \mu_{\Omega}^j(\mathbf{z}, \mathbf{w}) = i \sum_l a_l^j (|z_l|^2 - |w_l|^2) + 2k \sum_l a_l^j \bar{w}_l z_l.$$

Denote by $\Delta_{\alpha_1, \dots, \alpha_k}$ the $\binom{n}{k}$ $k \times k$ minor determinants of Ω .

Definition 5.11. Let $\Omega \in \mathcal{M}_{k,n}(\mathbb{Z})$ be a weight matrix.

(i) Ω is nondegenerate if $\Delta_{\alpha_1, \dots, \alpha_k} \neq 0$, for all $1 \leq \alpha_1 < \dots < \alpha_k \leq n$.

Let Ω be nondegenerate, and let d be the gcd of all the $\Delta_{\alpha_1, \dots, \alpha_k}$, the k th determinantal divisor. Then Ω is admissible if

(ii) $\gcd(\Delta_{\alpha_2, \dots, \alpha_{k+1}}, \dots, \Delta_{\alpha_1, \dots, \hat{\alpha}_t, \dots, \alpha_{k+1}}, \dots, \Delta_{\alpha_1, \dots, \alpha_k}) = d$ for all length $k+1$ sequences $1 \leq \alpha_1 < \dots < \alpha_t < \dots < \alpha_{k+1} \leq n+1$.

The quotient obtained in Proposition 5.10 $\mathcal{S}_\Omega = S^{4n-1} // T^k(\Omega)$ will depend on Ω only up to a certain equivalence. Choosing a different basis of $\mathfrak{t}_k$ results in an action on Ω by an element in $Gl(k, \mathbb{Z})$. We also have the normalizer of T^n in $Sp(n)$, the Weyl group $\mathcal{W}(Sp(n)) = \Sigma_n \times \mathbb{Z}_2^n$ where Σ_n is the permutation group. $\mathcal{W}(Sp(n))$ acts on S^{4n-1} preserving the 3-Sasakian structure, and it acts on weight matrices by permutations and sign changes of columns. The group $Gl(k, \mathbb{Z}) \times \mathcal{W}(Sp(n))$ acts on $\mathcal{M}_{k,n}(\mathbb{Z})$. We say Ω is *reduced* if $d = 1$ in definition 5.11. It is a result in [8] that we may assume that a nondegenerate weight matrix Ω is reduced, as this is the case when (5.4) is an inclusion.

Theorem 5.12 ([5, 8]). Let $\Omega \in \mathcal{M}_{k,n}(\mathbb{Z})$ be reduced.

(i) If Ω is nondegenerate, then $\mathcal{S}_\Omega$ is an orbifold.

(ii) Supposing Ω is nondegenerate, $\mathcal{S}_\Omega$ is smooth if and only if Ω is admissible.

Notice that the automorphism group of $\mathcal{S}_\Omega$ contains $T^{n-k} \cong T^n / \iota_\Omega(T^k)$.

We now restrict to the case of 7-dimensional toric quotients, so $n = k + 2$. We may take matrices of the form

$$(5.7) \quad \Omega = \begin{pmatrix} 1 & 0 & \cdots & 0 & a_1 & b_1 \\ 0 & 1 & \cdots & 0 & a_2 & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & a_k & b_k \end{pmatrix}.$$

Proposition 5.13 ([8]). Let $\Omega \in \mathcal{M}_{k,k+2}(\mathbb{Z})$ be as above. Then Ω is admissible if and only if $a_i, b_j, i, j = 1, \dots, k$ are all nonzero, $\gcd(a_i, b_i) = 1$ for $i = 1, \dots, k$, and we do not have $a_i = a_j$ and $b_i = b_j$, or $a_i = -a_j$ and $b_i = -b_j$ for some $i \neq j$.

This shows that for $n = k + 2$ there are infinitely many reduced admissible weight matrices. One can, for example, choose $a_i, b_j, i, j = 1, \dots, k$ be all pairwise relatively prime. We will make use of the cohomology computation of R. Hepworth [26] to show that we have infinitely many smooth 3-Sasakian 7-manifolds of each second Betti number $b_2 \geq 1$. Let $\Delta_{p,q}$ denote the $k \times k$ minor determinant of Ω obtained by deleting the p^{th} and q^{th} columns.

Theorem 5.14 ([26][7, 8]). Let $\Omega \in \mathcal{M}_{k,k+2}(\mathbb{Z})$ be a reduced admissible weight matrix. Then $\pi_1(\mathcal{S}_\Omega) = e$. And the cohomology of $\mathcal{S}_\Omega$ is

p	0	1	2	3	4	5	6	7
H^p	$\mathbb{Z}$	0	$\mathbb{Z}^k$	0	G_Ω	$\mathbb{Z}^k$	0	$\mathbb{Z}$

where G_Ω is a torsion group of order

$$\sum |\Delta_{s_1, t_1}| \cdots |\Delta_{s_{k+1}, t_{k+1}}|$$

with the summand with index $s_1, t_1, \dots, s_{k+1}, t_{k+1}$ included if and only if the graph on the vertices $\{1, \dots, k+2\}$ with edges $\{s_i, t_i\}$ is a tree.

If we consider weight matrices as in Proposition 5.13, then the order of G_Ω is greater than $|a_1 \cdots a_k| + |b_1 \cdots b_k|$. We have the following.

Corollary 5.15 ([26][8]). *There are smooth toric 3-Sasakian 7-manifolds with second Betti number $b_2 = k$ for all $k \geq 0$. Furthermore, there are infinitely many possible homotopy types of examples $\mathcal{S}_\Omega$ for each $k > 0$.*

5.3. Anti-self-dual Einstein spaces. We consider the anti-self-dual Einstein orbifolds $\mathcal{M} = \mathcal{M}_\Omega$ associated to the toric 3-Sasakian 7-manifolds $\mathcal{S}$ in greater detail. Since $\mathcal{M}$ is a 4 dimensional orbifold with an effective action of T^2 the techniques of [25] show that $\mathcal{M}$ is characterized by the polygon $\mathcal{Q}_\Omega = \mathcal{M}/T^2$ with $k+2$ edges, $b_2(\mathcal{M}) = k$, labeled in cyclic order with $(m_0, n_0), (m_1, n_1), \dots, (m_{k+2}, n_{k+2})$ in $\mathbb{Z}^2$, $(m_0, n_0) = -(m_{k+2}, n_{k+2})$, denoting the isotropy subgroups. For the quotients $\mathcal{M}_\Omega$ one can show the following (cf [40]):

- a. The sequence m_i , $i = 0, \dots, k+2$ is strictly increasing.
- b. The sequence $(n_i - n_{i-1})/(m_i - m_{i-1})$, $i = 1, \dots, k+2$ is strictly increasing.

We will make use of the following result of D. Calderbank and M. Singer [12] which classifies those compact orbifolds which admit toric anti-self-dual Einstein metrics. The case for which the associated 3-Sasakian space is smooth is originally due to R. Bielawski [4].

Theorem 5.16. *Let $\mathcal{M}$ be a compact toric 4-orbifold with $\pi_1^{orb}(\mathcal{M}) = e$ and $k = b_2(\mathcal{M})$. Then the following are equivalent.*

- (i) *One can arrange that the isotropy data of $\mathcal{M}$ satisfy a. and b. above by cyclic permutations, changing signs, and acting by $Gl(2, \mathbb{Z})$.*
- (ii) *$\mathcal{M}$ admits a toric anti-self-dual Einstein metric unique up to homothety and equivariant diffeomorphism. Furthermore, $(\mathcal{M}, g)$ is isometric to the quaternionic Kähler reduction of $\mathbb{H}P^{k+1}$ by a torus $T^k \subset Sp(k+2)$.*

It is well known that the only possible smooth compact anti-self-dual Einstein spaces with positive scalar curvature are S^4 and $\overline{\mathbb{C}P}^2$, which are both toric.

Suppose $\mathcal{M}$ has isotropy data $v_0, v_1, \dots, v_{k+2}$. Then it is immediate that $v_0, v_1, \dots, v_{k+2}, -v_1, -v_2, \dots, -v_{k+1}$ are the vertices of a convex polygon in $N_{\mathbb{R}} = \mathbb{R}^2$, which defines an augmented fan Δ^* defining a toric Fano surface X . A bit more thought gives the following.

Theorem 5.17. *There is a one to one correspondence between compact toric anti-self-dual Einstein orbifolds $\mathcal{M}$ with $\pi_1^{orb}(\mathcal{M}) = e$ and special symmetric toric Fano orbifold surfaces X with $\pi_1^{orb}(X) = e$. And X has a Kähler-Einstein metric of positive scalar curvature. Under the correspondence if $b_2(\mathcal{M}) = k$, then $b_2(X) = 2k + 2$.*

This will be reproved in section 5.4 by exhibiting X as a divisor in the twistor space.

Example. Consider the admissible weight matrix

$$\Omega = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 2 \end{pmatrix}.$$

28

Then the 3-Sasakian space $\mathcal{S}_\Omega$ is smooth and $b_2(\mathcal{S}_\Omega) = b_2(\mathcal{M}_\Omega) = 2$. And the anti-self-dual orbifold $\mathcal{M}_\Omega$ has isotropy data

$$v_0 = (-7, -2), (-5, -2), (-1, -1), (5, 1), (7, 2) = v_4.$$

The singular set of $\mathcal{M}$ consists of two points with stablizer group $\mathbb{Z}_3$ and two with $\mathbb{Z}_4$. The associated toric Kähler-Einstein surface is that in figure 2. $\diamond$

The generic fiber of $\varpi : \mathcal{S} \rightarrow \mathcal{M}$ is either $SO(3)$ or $Sp(1)$. In general the existence of a lifting to an $Sp(1)$ V-bundle is obstructed by the *Marchiafava-Romani class* which when $\mathcal{M}$ is 4-dimensional is identical to $w_2(\mathcal{M}) \in H_{orb}^2(\mathcal{M}, \mathbb{Z}_2)$. In other words, the contact line bundle $\mathbf{L}$ has a square root $\mathbf{L}^{\frac{1}{2}}$ if, and only if, $w_2(\mathcal{M}) = 0$.

Proposition 5.18. *Let X be the symmetric toric Fano surface associated to the anti-self-dual Einstein orbifold $\mathcal{M}$. Then $\text{Ind}(X) = 2$ if and only if $w_2(\mathcal{M}) = 0$. In other words, $\mathbf{K}_X^{-1}$ has a square root if and only if the contact line bundle $\mathbf{L}$ does.*

See [40] Proposition 5.22 for a proof.

5.4. Twistor space and divisors. We will consider the twistor space $\mathcal{Z}$ introduced in Theorem 5.6 more closely for the case when $\mathcal{M}$ is an anti-self-dual Einstein orbifold. For now suppose $(\mathcal{M}, [g])$ is an anti-self-dual, i.e. $W_+ \equiv 0$, conformal orbifold. There is a twistor space of $(\mathcal{M}, [g])$ which is originally due to R. Penrose [36]. See [1] for positive definite case.

The *twistor space* of $(\mathcal{M}, [g])$ is a complex three dimensional orbifold $\mathcal{Z}$ with the following properties:

- a. There is a V-bundle fibration $\varpi : \mathcal{Z} \rightarrow \mathcal{M}$.
- b. The general fiber of $P_x = \varpi^{-1}(x), x \in \mathcal{Z}$ is a projective line $\mathbb{CP}^1$ with normal bundle $N \cong \mathcal{O}(1) \oplus \mathcal{O}(1)$, which holds over singular fibers with N a V-bundle.
- c. There exists an anti-holomorphic involution σ of $\mathcal{Z}$ leaving the fibers P_x invariant.

Let T be an oriented real 4-dimensional vector space with inner product g . Let $C(T)$ be set of orthogonal complex structures inducing the orientation, i.e. if $r, s \in T$ is a complex basis then r, Jr, s, Js defines the orientation. One has $C(T) = S^2 \subset \Lambda_+^2(T)$, where S^2 is the sphere of radius $\sqrt{2}$. Now take T to be $\mathbb{H}$. Recall that $Sp(1)$ is the group of unit quaternions. Let

$$(5.8) \quad Sp(1)_+ \times Sp(1)_-$$

act on $\mathbb{H}$ by

$$(5.9) \quad w \rightarrow gw g'^{-1}, \text{ for } w \in \mathbb{H} \text{ and } (g, g') \in Sp(1)_+ \times Sp(1)_-.$$

Then we have

$$(5.10) \quad Sp(1)_+ \times_{\mathbb{Z}_2} Sp(1)_- \cong SO(4),$$

where $\mathbb{Z}_2$ is generated by $(-1, -1)$. Let

$$(5.11) \quad \begin{aligned} C &= \{ai + bj + ck : a^2 + b^2 + c^2 = 1, a, b, c \in \mathbb{R}\} \\ &= \{g \in Sp(1)_+ : g^2 = -1\} \cong S^2. \end{aligned}$$

Then $g \in C$ defines an orthogonal complex structure by

$$w \rightarrow gw, \text{ for } w \in \mathbb{H},$$

giving an identification $C = C(\mathbb{H})$. Let $V_+ = \mathbb{H}$ considered as a representation of $Sp(1)_+$ and a right $\mathbb{C}$ -vector space. Define $\pi : V_+ \setminus \{0\} \rightarrow C$ by $\pi(h) = -h i h^{-1}$. Then the fiber of π over $h i h^{-1}$ is $h\mathbb{C}$. Then π is equivariant if $Sp(1)_+$ acts on C by $q \rightarrow g q g^{-1}, g \in Sp(1)_+$. We have a the identification

$$(5.12) \quad C = V_+ \setminus \{0\} / \mathbb{C}^* = \mathbb{P}(V_+).$$

Fix a Riemannian metric g in $[g]$. Let $\phi : \tilde{U} \rightarrow U \subset \mathcal{M}$ be a local uniformizing chart with group Γ . Let $F_{\tilde{U}}$ be the bundle of orthonormal frames on $\tilde{U}$. Then

$$(5.13) \quad F_{\tilde{U}} \times_{SO(4)} \mathbb{P}(V_+) = F_{\tilde{U}} \times_{SO(4)} C$$

defines a local uniformizing chart for $\mathcal{Z}$ mapping to

$$F_{\tilde{U}} \times_{SO(4)} \mathbb{P}(V_+) / \Gamma = F_{\tilde{U}} / \Gamma \times_{SO(4)} \mathbb{P}(V_+).$$

Right multiplication by j on $V_+ = \mathbb{H}$ defines the anti-holomorphic involution σ which is fixed point free on (5.13). We will denote a neighborhood as in (5.13) by $\tilde{U}_{\mathcal{Z}}$.

An almost complex structure is defined as follows. At a point $z \in \tilde{U}_{\mathcal{Z}}$ the Levi-Civita connection defines a horizontal subspace H_z of the real tangent space T_z and we have a splitting

$$(5.14) \quad T_z = H_z \oplus T_z P_x = T_x \oplus T_z P_x,$$

where $\varpi(z) = x$ and T_x is the real tangent space of $\tilde{U}$. Let J_z be the complex structure on T_x given by $z \in P_x = C(T_x)$, and let J'_z be complex structure on $T_x \oplus T_z P_x$ arising from the natural complex structure on P_x . Then the almost complex structure on T_z is the direct sum of J_z and J'_z . This defines a natural almost complex structure on $Z_{\tilde{U}}$ which is invariant under Γ . We get an almost complex structure on $\mathcal{Z}$ which is integrable precisely when $W_+ \equiv 0$.

Assume that $\mathcal{M}$ anti-self-dual Einstein with non-zero scalar curvature. Then $\mathcal{Z}$ has a complex contact structure $D \subset T^{1,0}\mathcal{Z}$ with holomorphic contact form $\theta \in \Gamma(\Lambda^{1,0}\mathcal{Z} \otimes \mathbf{L})$ where $\mathbf{L} = T^{1,0}\mathcal{Z}/D$.

The group of isometries $\text{Isom}(\mathcal{M})$ lifts to an action on $\mathcal{Z}$ by real holomorphic transformations. Real means commuting with σ . This extends to a holomorphic action of the complexification $\text{Isom}(\mathcal{M})_{\mathbb{C}}$. For $X \in \mathfrak{Isom}(\mathcal{M}) \otimes \mathbb{C}$, the Lie algebra of $\text{Isom}(\mathcal{M})_{\mathbb{C}}$, we will also denote by X the holomorphic vector field induced on $\mathcal{Z}$. Then $\theta(X) \in H^0(\mathcal{Z}, \mathcal{O}(\mathbf{L}))$. By a well known twistor correspondence the map $X \rightarrow \theta(X)$ defines an isomorphism

$$(5.15) \quad \mathfrak{Isom}(\mathcal{M}) \otimes \mathbb{C} \cong H^0(\mathcal{Z}, \mathcal{O}(\mathbf{L})),$$

which maps real vector fields to real sections of $\mathbf{L}$.

Suppose for now on that $\mathcal{M}$ is a toric anti-self-dual Einstein orbifold with twistor space $\mathcal{Z}$. We will assume that $\pi_1^{orb}(\mathcal{M}) = e$ which can always be arranged by taking the orbifold cover. Then as above T^2 acts on $\mathcal{Z}$ by holomorphic transformations. And the action extends to $T_{\mathbb{C}}^2 = \mathbb{C}^* \times \mathbb{C}^*$, which in this case is an algebraic action. Let $\mathfrak{t}$ be the Lie algebra of T^2 with $\mathfrak{t}_{\mathbb{C}}$ the Lie algebra of $T_{\mathbb{C}}^2$. Then we have from (5.15) the pencil

$$(5.16) \quad E = \mathbb{P}(\mathfrak{t}_{\mathbb{C}}) \subseteq |\mathbf{L}|,$$

where for $t \in E$ we denote $X_t = (\theta(t))$ the divisor of the section $\theta(t) \in H^0(\mathcal{Z}, \mathcal{O}(\mathbf{L}))$. Note that E has an equator of real divisors. Also, since $T_{\mathbb{C}}^2$ is abelian, every $X_t, t \in E$ is $T_{\mathbb{C}}^2$ invariant.

Our goal is to determine the structure of the divisors in the pencil E . As before we will consider the one parameter groups $\rho_i \in N = \mathbb{Z} \times \mathbb{Z}$, where N is the lattice of one parameter $\mathbb{C}^*$ -subgroups of $T_{\mathbb{C}}^2$. Also, we will identify the Lie algebra $\mathfrak{t}$ of T^2 with $N \otimes \mathbb{R}$ and the Lie algebra $\mathfrak{t}_{\mathbb{C}}$ of $T_{\mathbb{C}}^2$ with $N \otimes \mathbb{C}$. Since $\mathbf{L}|_{P_x} = \mathcal{O}(2)$ a divisor $X_t \in E$ intersects a generic twistor line P_x at two points.

Recall that the set of non-trivial stablizers of the T^2 -action on $\mathcal{M}$ is $B = \cup_{i=1}^{k+2} B_i$ where B_i is topologically a 2-sphere. Denote by $x_i = B_i \cap B_{i+1}$ the $k+2$ fixed points of the action. We will denote $P_i := P_{x_i}, i = 1, \dots, k+2$. One can show there exist two irreducible rational curves $C_i^{\pm}, i = 1, \dots, k+2$ mapped diffeomorphically to B_i by ϖ . Furthermore, $\sigma(C_i^{\pm}) = C_i^{\mp}$. The singular set for the T^2 -action on $\mathcal{Z}$ is the union of rational curves

$$(5.17) \quad \Sigma = \left(\cup_{i=1}^{k+2} P_i \right) \cup \left(\cup_{i=1}^{k+2} C_i^+ \cup C_i^- \right).$$

With a closer analysis of the action of $T_{\mathbb{C}}^2$ on $\mathcal{Z}$ one can prove the following. See [40] for the proof. The term *suborbifold* denotes a subvariety which is a submanifold in every local uniformizing neighborhood.

Theorem 5.19. *Let $\mathcal{M}$ be a compact anti-self-dual Einstein orbifold with $b_2(\mathcal{M}) = k$ and $\pi_1^{orb}(\mathcal{M}) = e$. Let $n = k + 2$. Then there are distinct real points $t_1, t_2, \dots, t_n \in E$ so that for $t \in E \setminus \{t_1, t_2, \dots, t_n\}$, $X_t \subset \mathcal{Z}$ is a suborbifold. Furthermore X_t is a special symmetric toric Fano surface. The anti-canonical cycle of X_t is $C_1, C_2, \dots, C_{2n}$, and the corresponding stabilizers are $\rho_1, \rho_2, \dots, \rho_{2n}$ which define the vertices in $N = \mathbb{Z} \times \mathbb{Z}$ of Δ^* with $X_t = X_{\Delta^*}$.*

For $t_i \in E$, $X_{t_i} = D + \bar{D}$, where $D, \bar{D}$ are irreducible degree one divisors with $\sigma(D) = \bar{D}$. The $D, \bar{D}$ are suborbifolds of $\mathcal{Z}$ and are toric Fano surfaces. We have $D \cap \bar{D} = P_i$ and the elements $\pm(\rho_1, \dots, \rho_i, -\rho_i + \rho_{i+1}, \rho_{n+i+1}, \dots, \rho_{2n})$ define the augmented fans for D and $\bar{D}$.

Note that a consequence of the theorem is that Σ given by equation (5.17) is the set with non-trivial stablizers of the action of the complex torus $T_{\mathbb{C}}^2$.

6. SASAKIAN SUBMANIFOLDS

In this section we use the results on toric 3-Sasakian manifolds to produce a new infinite series of toric Sasaki-Einstein 5-manifolds corresponding to the toric 3-Sasakian 7-manifolds discussed above and completing the correspondence in diagram 1.1. But first we review the Smale/Barden classification of smooth 5-manifolds which will be used. The possible diffeotypes of toric examples in dimension 5 is very limited.

6.1. Classification of 5-Manifolds. Closed smooth simply connected spin 5-manifolds were classified by S. Smale [38]. Subsequently D. Barden extended the classification to the non-spin case [3]. Consider the primary decomposition

$$(6.1) \quad H_2(M, \mathbb{Z}) \cong \mathbb{Z}^r \oplus \mathbb{Z}_{k_1} \oplus \mathbb{Z}_{k_2} \oplus \dots \oplus \mathbb{Z}_{k_s},$$

where k_j divides k_{j+1} . Of course the decomposition is not unique, but the $r, k_1, \dots, k_s$ are. The second Stiefel-Whitney class defines a homomorphism $w_2 : H_2(M, \mathbb{Z}) \rightarrow \mathbb{Z}_2$. One can arrange the decomposition 6.1 so that w_2 is non-zero on only one component $\mathbb{Z}_{k_j}$, or $\mathbb{Z}$ of $\mathbb{Z}^r$. Then define $i(M)$ to be i if 2^i is the 2-primary component of $\mathbb{Z}_{k_j}$, or ∞ . Alternatively, $i(M)$ is the minimum i so that w_2 is non-zero on a 2-primary component of order 2^i of $H_2(M, \mathbb{Z})$.

Theorem 6.1 ([38, 3]). *Smooth simply connected closed 5-manifolds are classifiable up to diffeomorphism. Any such manifold is diffeomorphic to one of*

$$X_j \# M_{k_1} \# \cdots \# M_{k_s},$$

where $-1 \leq j \leq \infty, s \geq 0, 1 < k_1$ and K_i divides k_{i+1} or $k_{i+1} = \infty$. A complete set of invariants is given by $H_2(M, \mathbb{Z})$ and $i(M)$, and the manifolds $X_{-1}, X_0, X_j, X_\infty, M_k, M_\infty$ are as follows

M	$H_2(M, \mathbb{Z})$	$i(M)$
$M_0 = X_0 = S^5$	0	0
$M_k, 1 < k < \infty$	$\mathbb{Z}_k \oplus \mathbb{Z}_k$	0
$M_\infty = S^2 \times S^3$	$\mathbb{Z}$	0
X_{-1}	$\mathbb{Z}_2$	1
$X_j, 0 < j < \infty$	$\mathbb{Z}_{2^j} \oplus \mathbb{Z}_{2^j}$	j
X_∞	$\mathbb{Z}$	∞

where $X_{-1} = SU(3)/SO(3)$ is the Wu manifold, and X_∞ is the non-trivial S^3 -bundle over S^2 .

The existence of an effective T^3 action severely restricts the topology by the following theorem of Oh [35].

Theorem 6.2. *Let M be a compact simply connected 5-manifold with an effective T^3 -action. Then $H_2(M, \mathbb{Z})$ has no torsion. Thus, M is diffeomorphic to S^5 , $\#kM_\infty$, or $X_\infty \# (k-1)M_\infty$, where $k = b_2(M) \geq 1$. Conversely, these manifolds admit effective T^3 -actions.*

By direct construction C. Boyer, K. Galicki, and L. Ornea [9] showed that the manifolds in this theorem admit toric Sasakian structures and, in fact, admit regular Sasakian structures.

A simply connected Sasaki-Einstein manifold must have $w_2 = 0$, therefore we have the following:

Corollary 6.3. *Let M be a compact simply connected 5-manifolds with a toric Sasaki-Einstein structure. Then M is diffeomorphic to S^5 , or $\#kM_\infty$, where $k = b_2(M) \geq 1$.*

6.2. Sasakian Submanifolds and Examples. Associated to each compact toric anti-self-dual Einstein orbifold $\mathcal{M}$ with $\pi_1^{orb}(\mathcal{M}) = e$ is the twistor space $\mathcal{Z}$ and a family of embeddings $X_t \subset \mathcal{Z}$ where $t \in E \setminus \{t_1, t_2, \dots, t_{k+2}\}$ and $X = X_t$ is the symmetric toric Fano surface canonically associated to $\mathcal{M}$. We denote the family of embeddings by

$$(6.2) \quad \iota_t : X \rightarrow \mathcal{Z}.$$

Let M be the total space of the S^1 V-bundle associated to $\mathbf{K}_X$ or $\mathbf{K}_X^{\frac{1}{2}}$, depending on whether $\text{Ind}(X) = 1$ or 2 .

Theorem 6.4. *Let $\mathcal{M}$ be a compact toric anti-self-dual Einstein orbifold with $\pi_1^{orb}(\mathcal{M}) = e$. There exists a Sasakian structure $\{\tilde{g}, \tilde{\Phi}, \tilde{\xi}, \tilde{\eta}\}$ on M , such that if $(X, \hat{h})$ is the Kähler structure making $\pi : M \rightarrow X$ a Riemannian submersion, then*

we have the following diagram where the horizontal maps are isometric embeddings.

$$(6.3) \quad \begin{array}{ccc} M & \xrightarrow{\bar{\iota}_t} & \mathcal{S} \\ \downarrow & & \downarrow \\ X & \xrightarrow{\iota_t} & \mathcal{Z} \\ & & \downarrow \\ & & \mathcal{M} \end{array}$$

If the 3-Sasakian space $\mathcal{S}$ is smooth, then so is M . If M is smooth, then

$$M \underset{\text{diff}}{\cong} \#kM_\infty, \text{ where } k = 2b_2(\mathcal{S}) + 1.$$

Proof. Let $\{g, \Phi, \xi, \eta\}$ be the fixed Sasakian structure on $\mathcal{S}$ with Φ descending to the complex structure on $\mathcal{Z}$. The adjunction formula gives

$$(6.4) \quad \mathbf{K}_X \cong \mathbf{K}_{\mathcal{Z}} \otimes [X]|_X = \mathbf{K}_{\mathcal{Z}} \otimes \mathbf{K}_{\mathcal{Z}}^{-\frac{1}{2}}|_X = \mathbf{K}_{\mathcal{Z}}^{\frac{1}{2}}|_X.$$

Let h be the Kähler-Einstein metric on $\mathcal{Z}$ related to the 3-Sasakian metric g on $\mathcal{S}$ by Riemannian submersion. Recall that $\mathcal{S}$ is the total space of the S^1 V-bundle associated to $\mathbf{L}^{-1}$, (resp. $\mathbf{L}^{-\frac{1}{2}}$ if $w_2(\mathcal{M}) = 0$). Also M is the total space of the S^1 V-bundle associated to $\mathbf{K}_X$, (resp. $\mathbf{K}_X^{\frac{1}{2}}$ if $w_2(\mathcal{M}) = 0$). Using the isomorphism in (6.4) we lift ι_t to $\bar{\iota}_t$. The metric on $\mathcal{S}$ is

$$g = \eta \otimes \eta + \pi^*h.$$

Pull back h and η to $\tilde{h} = \iota_t^*h$ and $\tilde{\eta} = \bar{\iota}_t^*\eta$ respectively. It follows that $\frac{1}{2}\tilde{\eta} = \tilde{\omega}$, where $\tilde{\omega}$ is the Kähler form of $\tilde{h}$. Then $\tilde{\Phi} = \tilde{\nabla}\tilde{\xi}$ is a lift of the complex structure J on X . And the integrability condition in definition 2.1 follows from the integrability of J . Then

$$\tilde{g} = \tilde{\eta} \otimes \tilde{\eta} + \pi^*\tilde{h}$$

is a Sasakian metric on M .

If $\mathcal{S}$ is smooth, then the orbifold uniformizing groups act on $\mathbf{L}^{-1}$ (or $\mathbf{L}^{\frac{1}{2}}$) minus the zero section without non-trivial stabilizers. By (6.4) this holds for the bundle $\mathbf{K}_X$ (or $\mathbf{K}_X^{\frac{1}{2}}$) on X .

It follows from $\pi_1^{\text{orb}}(\mathcal{M}) = e$ that $\pi_1^{\text{orb}}(X) = e$ (cf. [25]). And $\pi_1^{\text{orb}}(M) = e$ by arguments as after Proposition 2.4. If M is smooth, Corollary 6.3 gives the diffeomorphism. $\square$

We are more interested in M with the Sasaki-Einstein metric that exists by corollary 4.6. In this case the horizontal maps are not isometries.

Consider the reducible cases, $t_i \in E, i = 1, \dots, k+2$, where $X_{t_i} = D + \bar{D} \subset \mathcal{Z}$. Then restricting $\mathbf{L}^{-1}$, (resp. $\mathbf{L}^{-\frac{1}{2}}$ if $w_2(\mathcal{M}) = 0$), and arguing as above we obtain Sasakian manifolds $N_i, \bar{N}_i \subset \mathcal{S}$, where smoothness follows from that of $\mathcal{S}$, whose Sasakian structures $(N_i, g_i, \Phi_i, \xi_i, \eta_i)$ and $(\bar{N}_i, \bar{g}_i, \bar{\Phi}_i, \bar{\xi}_i, \bar{\eta}_i)$ pull back from that of $\mathcal{S}$. And $N_{t_i} \cap \bar{N}_{t_i}$ is a lens space with the constant curvature metric. Note that ς restricts to an isometry $\varsigma : N_i \rightarrow \bar{N}_i$ which gives a conjugate isomorphism between Sasakian structures. These manifolds do not satisfy the conditions of Proposition 4.1, so cannot be transversally deformed to Sasaki-Einstein structures. But $c_1^B(N_i) > 0$, so they can be transversally deformed to positive Ricci curvature Sasakian by the transverse Calabi-Yau theorem [28]. By Theorem 6.2 N_i is diffeomorphic to $\#kM_\infty$, or $X_\infty \#(k-1)M_\infty$, where $k = b_2(N_i) = b_2(\mathcal{S})$. By the remarks

after Proposition 7.1 $N_i, i = 1, \dots, k+2$, is diffeomorphic to $X_\infty \# (k-1)M_\infty$ if $w_2(\mathcal{M}) \neq 0$.

The family of submanifolds $\bar{\iota}_t : M \rightarrow \mathcal{S}$ for $t \in E \setminus \{t_1, t_2, \dots, t_{k+2}\}$ and $N_i, \bar{N}_i, i = 1, \dots, k+2$ for the reducible cases have a simple description. Recall the 1-form $\eta = \eta^2 + i\eta^3$ of section 5.1 which is $(1, 0)$ with respect to the CR structure $I = -\Phi^1$. For $t \in \mathfrak{t}$ let Y_t denote the killing vector field on $\mathcal{Z}$ with lift $\bar{Y}_t \in I(\mathcal{S}, g)$. Then $\theta(Y_t) \in H^0(\mathcal{Z}, \mathcal{O}(\mathbf{L}))$ which defines a holomorphic function on $\mathbf{L}^{-1}$. Since $\mathcal{S}$ is the S^1 subbundle of $\mathbf{L}^{-1}$, we have $\theta(Y_t) = \eta(\bar{Y}_t)$. Complexifying gives the same equality for $t \in \mathfrak{t}_\mathbb{C}$. Thus for $t \in E \setminus \{t_1, t_2, \dots, t_{k+2}\}$, we have $M_t := \bar{\iota}_t(M) = (\eta(\bar{Y}_t)) \subset \mathcal{S}$ and $N_i \cup \bar{N}_i = (\eta(\bar{Y}_{t_i})) \subset \mathcal{S}$, where of course $(\eta(\bar{Y}_t))$ denotes the submanifold $\eta(\bar{Y}_t) = 0$. Note that here we are setting $2/3$ s of the moment map to zero.

This gives us the new infinite families of Sasaki-Einstein manifolds and the diagram 1.1.

Theorem 6.5. *Let $(\mathcal{S}, g)$ be a toric 3-Sasakian 7-manifold with $\pi_1(\mathcal{S}) = e$. Canonically associated to $(\mathcal{S}, g)$ are a special symmetric toric Fano surface X and a toric quasi-regular Sasaki-Einstein 5-manifold (M, g, Φ, ξ, η) which fit in the commutative diagram 6.3. We have $\pi_1^{orb}(X) = e$ and $\pi_1(M) = e$. And*

$$M \underset{\text{diff}}{\cong} \#kM_\infty, \text{ where } k = 2b_2(\mathcal{S}) + 1$$

Furthermore $(\mathcal{S}, g)$ can be recovered from either X or M with their torus actions.

Proof. Note that the homotopy sequence

$$\cdots \rightarrow \pi_1(G) \rightarrow \pi_1(\mathcal{S}) \rightarrow \pi_1^{orb}(\mathcal{M}) \rightarrow e,$$

where $G = SO(3)$ or $Sp(1)$, shows that $\pi_1^{orb}(\mathcal{M}) = e$. The toric surface X_{Δ^*} and Sasakian 5-manifold $(M, \tilde{g}, \tilde{\Phi}, \tilde{\xi}, \tilde{\eta})$ are given in Theorem 6.4. By Corollary 4.6 this Sasakian structure has a transversal deformation to a Sasaki-Einstein structure.

Given either X or M , by the discussion in section 5.3, the orbifold $\mathcal{M}$ can be recovered. By Theorem 5.16 $\mathcal{M}$ admits a unique anti-self-dual Einstein structure up to homothety compatible with the torus action. The 3-Sasakian space $\mathcal{S}$ and its twistor space $\mathcal{Z}$ can be constructed from $\mathcal{M}$ and its anti-self-dual Einstein structure. (cf. [5]). $\square$

Corollary 6.6. *For each odd $k \geq 3$ there is a countably infinite number of distinct toric quasi-regular Sasaki-Einstein structures on $\#kM_\infty$.*

We do not know the Sasaki-Einstein metrics explicitly. But if $c = \text{Ind}(M)$, an application of Corollary 3.27 gives

$$\begin{aligned} \text{Vol}(M, g) &= \frac{2\pi c}{3} \text{Vol}(X, \omega) \\ &= 2c \left(\frac{\pi}{3}\right)^3 \text{Vol}(\Sigma_{-k}), \end{aligned}$$

where ω is the transversal Kähler metric. We have $c = 1$ or 2 . Let $(M, g_i), i \in \mathbb{Z}_+$, be any infinite sequence of metrics on $\#kM_\infty$ in Corollary 6.6. These Sasaki-Einstein structures have leaf spaces X_i , where $X_i = X_{\Delta_i^*}$. Observe that the polygons Δ_i^* get arbitrarily large, and the anti-canonical polytopes $(\Sigma_{-k})_i$ satisfy

$$\text{Vol}((\Sigma_{-k})_i) \rightarrow 0, \text{ as } i \rightarrow \infty.$$

Thus we have $\text{Vol}(M, g_i) \rightarrow 0$, as $i \rightarrow \infty$.

6.3. Examples. We consider some of the examples obtained starting with the simplest. In particular we can determine some of the spaces in diagram 1.1 associated to a toric 3-Sasakian 7-manifold more explicitly in some cases.

6.3.1. Smooth examples. It is well known that there exists only two complete examples of positive scalar curvature anti-self-dual Einstein manifolds [27] [15], S^4 and $\mathbb{CP}^2$ with the round and Fubini-Study metrics respectively. Note that we are considering $\mathbb{CP}^2$ with the opposite of the usual orientation.

$$\mathcal{M} = S^4$$

For the spaces in diagram 1.1 we have: $\mathcal{M} = S^4$ with the round metric; its twistor space $\mathcal{Z} = \mathbb{CP}^3$ with the Fubini-study metric; the quadratic divisor $X \subset \mathcal{Z}$ is $\mathbb{CP}^1 \times \mathbb{CP}^1$ with the homogeneous Kähler-Einstein metric; $M = S^2 \times S^3$ with the homogeneous Sasaki-Einstein structure; and $\mathcal{S} = S^7$ has the round metric. In this case diagram 1.1 becomes the following.

$$(6.5) \quad \begin{array}{ccc} S^2 \times S^3 & \longrightarrow & S^7 \\ \downarrow & & \downarrow \\ \mathbb{CP}^1 \times \mathbb{CP}^1 & \longrightarrow & \mathbb{CP}^3 \\ & & \downarrow \\ & & S^4 \end{array}$$

This is the only example, I am aware of, for which the horizontal maps are isometric immersions when the toric surface and Sasakian space are equipped with the Einstein metrics.

$$\mathcal{M} = \mathbb{CP}^2$$

In this case $\mathcal{M} = \mathbb{CP}^2$ with the Fubini-Study metric and the reverse of the usual orientation; its twistor space is $\mathcal{Z} = F_{1,2}$, the manifold of flags $V \subset W \subset \mathbb{C}^3$ with $\dim V = 1$ and $\dim W = 2$, with the homogeneous Kähler-Einstein metric. The projection $\pi : F_{1,2} \rightarrow \mathbb{CP}^2$ is as follows. If $(p, l) \in F_{1,2}$ so l is a line in $\mathbb{CP}^2$ and $p \in l$, then $\pi(p, l) = p^\perp \cap l$, where $p^\perp$ is the orthogonal complement with respect to the standard hermitian inner product. We can define $F_{1,2} \subset \mathbb{CP}^2 \times (\mathbb{CP}^2)^*$ by

$$F_{1,2} = \{([p_0 : p_1 : p_2], [q^0 : q^1 : q^2]) \in \mathbb{CP}^2 \times (\mathbb{CP}^2)^* : \sum p_i q^i = 0\}.$$

And the complex contact structure is given by $\theta = q^i dp_i - p_i dq^i$. Fix the action of T^2 on $\mathbb{CP}^2$ by

$$(e^{i\theta}, e^{i\phi})[z_0 : z_1 : z_2] = [z_0 : e^{i\theta} z_1 : e^{i\phi} z_2].$$

Then this induces the action on $F_{1,2}$

$$(e^{i\theta}, e^{i\phi})([p_0 : p_1 : p_2], [q^0 : q^1 : q^2]) = ([p_0 : e^{i\theta} p_1 : e^{i\phi} p_2], [q^0 : e^{-i\theta} q^1 : e^{-i\phi} q^2]).$$

Given $[a, b] \in \mathbb{CP}^1$ the one parameter group $(e^{ia\tau}, e^{ib\tau})$ induces the holomorphic vector field $W_\tau \in \Gamma(T^{1,0}F_{1,2})$ and the quadratic divisor $X_\tau = (\theta(W_\tau))$ given by

$$X_\tau = (ap_1 q^1 + bp_2 q^2 = 0, \quad p_i q^i = 0).$$

One can check directly that X_τ is smooth for $\tau \in \mathbb{CP}^1 \setminus \{[1, 0], [0, 1], [1, 1]\}$ and $X_\tau = \mathbb{CP}_{(3)}^2$, the equivariant blow-up of $\mathbb{CP}^2$ at 3 points. For $\tau \in \{[1, 0], [0, 1], [1, 1]\}$, $X_\tau = D_\tau + \bar{D}_\tau$ where both $D_\tau, \bar{D}_\tau$ are isomorphic to the Hirzebruch surface $F_1 = \mathbb{P}(\mathcal{O}_{\mathbb{CP}^1} \oplus \mathcal{O}_{\mathbb{CP}^1}(1))$.

The Sasaki-Einstein space is $M = \#3(S^2 \times S^3)$. And the Sasakian manifolds $N_\tau, \bar{N}_\tau$ are diffeomorphic to X_∞ . $\mathcal{S} = \mathcal{S}(1, 1, 1) = SU(3)/U(1)$ with the homogeneous 3-Sasakian structure. This case has the following diagram.

$$(6.6) \quad \begin{array}{ccc} \#3(S^2 \times S^3) & \longrightarrow & SU(3)/U(1) \\ \downarrow & & \downarrow \\ \mathbb{C}P_{(3)}^2 & \longrightarrow & F_{1,2} \\ & & \downarrow \\ & & \mathbb{C}P^2 \end{array}$$

6.3.2. Galicki-Lawson quotients. The simplest examples of quaternionic-Kähler quotients are the Galicki-Lawson examples first appearing in [19] and further considered in [8]. These are circle quotients of $\mathbb{H}P^2$. In this case the weight matrices are of the form $\Omega = \mathbf{p} = (p_1, p_2, p_3)$ with the admissible set

$$\{\mathcal{A}_{1,3}(Z) = \{\mathbf{p} \in \mathbb{Z}^3 | p_i \neq 0 \text{ for } i = 1, 2, 3 \text{ and } \gcd(p_i, p_j) = 1 \text{ for } i \neq j\}$$

We may take $p_i > 0$ for $i = 1, 2, 3$. The zero locus of the 3-Sasakian moment map $N(\mathbf{p}) \subset S^{11}$ is diffeomorphic to the Stiefel manifold $V_{2,3}^{\mathbb{C}}$ of complex 2-frames in $\mathbb{C}^3$ which can be identified as $V_{2,3}^{\mathbb{C}} \cong U(3)/U(1) \cong SU(3)$. Let $f_{\mathbf{p}} : U(1) \rightarrow U(3)$ be

$$f_{\mathbf{p}}(\tau) = \begin{bmatrix} \tau^{p_1} & 0 & 0 \\ 0 & \tau^{p_2} & 0 \\ 0 & 0 & \tau^{p_3} \end{bmatrix}.$$

Then the 3-Sasakian space $\mathcal{S}(\mathbf{p})$ is diffeomorphic to the quotient of $SU(3)$ by the action of $U(1)$

$$\tau \cdot W = f_{\mathbf{p}}(\tau) W f_{(0,0,-p_1-p_2-p_3)}(\tau) \text{ where } \tau \in U(1) \text{ and } W \in SU(3).$$

Thus $\mathcal{S}(\mathbf{p}) \cong SU(3)/U(1)$ is a biquotient similar to the examples considered by Eschenburg in [14].

The action of the group $SU(2)$ generated by $\{\xi^1, \xi^2, \xi^3\}$ on $N(\mathbf{p}) \cong SU(3)$ commutes with the action of $U(1)$. We have $N(\mathbf{p})/SU(2) \cong SU(3)/SU(2) \cong S^5$ with $U(1)$ acting by

$$\tau \cdot v = f_{(-p_2-p_3, -p_1-p_3, -p_1-p_2)}(\tau) v \text{ for } v \in S^5 \subset \mathbb{C}^3.$$

We see that $\mathcal{M}_\Omega \cong \mathbb{C}P_{a_1, a_2, a_3}^2$ where $a_1 = p_2 + p_3, a_2 = p_1 + p_3, a_3 = p_1 + p_2$ and the quotient metric is anti-self-dual with the reverse of the usual orientation. If p_1, p_2, p_3 are all odd then the generic leaf of the 3-Sasakian foliation $\mathcal{F}_3$ is $SO(3)$. If exactly one is even, then the generic leaf is $Sp(1)$. Denote by X_{p_1, p_2, p_3} the toric Fano divisor, which can be considered as a generalization of $\mathbb{C}P_{(3)}^2$. We have the following spaces and embeddings.

$$(6.7) \quad \begin{array}{ccc} \#3(S^2 \times S^3) & \longrightarrow & \mathcal{S}(p_1, p_2, p_3) \\ \downarrow & & \downarrow \\ X_{p_1, p_2, p_3} & \longrightarrow & \mathcal{Z}(p_1, p_2, p_3) \\ & & \downarrow \\ & & \mathbb{C}P_{a_1, a_2, a_3}^2 \end{array}$$

7. POSITIVE RICCI CURVATURE EXAMPLES

In this section we use the toric geometry developed to construct examples of positive Ricci curvature Sasakian structures on the manifolds $X_\infty \# (k-1)M_\infty$ in Theorem 6.1. By Theorem 6.2 these are the only simply connected non-spin 5-manifolds that can admit toric Sasakian structures. They are already known to admit Sasakian structures [9].

Define a marked fan $\Delta^* = \Delta_{k,p}^*, k \geq 2, p \geq 0$ as follows. Let $\sigma_0 = (-1, 0)$, $\sigma_1 = (0, 1)$; $\sigma_j = (j-1, \frac{j(j-1)}{2} - 1), j = 2, \dots, k$; $\sigma_{k+1} = (k, \frac{(k+1)k}{2} - 1 + p)$, $\sigma_{k+2} = (0, \frac{(k+1)k}{2} + p)$. For $k = 1$ define $\Delta_{1,p}^*, p \geq 0$ by $\sigma_0 = (-1, 0), \sigma_1 = (0, 1), \sigma_2 = (1, 1 + p), \sigma_3 = (0, 2 + p)$. And define $l \in \text{SF}(\Delta^*)$ by

$$l(\sigma_j) := \begin{cases} 0 & \text{if } j = 0, \\ -1 & \text{if } 1 \leq j \leq k+2. \end{cases}$$

It is easy to check that l is strictly upper convex, and we have $\pi_1^{orb}(X) = e$ for all of the above fans. The cone $\mathcal{C}_l$ corresponding to $(\mathbf{L}_l^{-1})^\times$ on X_{Δ^*} satisfies the smoothness condition of equation 3.4, and $(\mathbf{L}_l^{-1})^\times$ is simply connected. Also, $-k \in \text{SF}(\Delta^*)$ is strictly upper convex. Thus by Corollary 3.25 $2\pi c_1(X_{\Delta^*}) = [\omega]$ for a Kähler form ω .

Proposition 7.1. *Each of the manifolds $M_{k,p}$, for $k \geq 1, p \geq 0$, is diffeomorphic to $X_\infty \# (k-1)M_\infty$. And for each p has a distinct positive Ricci curvature Sasakian structure.*

Proof. Let $X = X_{\Delta_{k,p}^*}$, and let $\mathbf{L}$ be the holomorphic line V-bundle associated with $l \in \text{SF}(\Delta^*)$. Since l is strictly upper convex, if ω is the canonical metric on X of Σ_l^* , then $[\omega] \in 2\pi c_1(\mathbf{L})$ by Theorem 3.24. By Proposition 2.4 there is a Sasakian structure (M, g, Φ, ξ, η) where M is the principle S^1 subbundle of $\mathbf{L}^{-1}$. The the toric cone $\mathcal{C}$ of $(\mathbf{L}^{-1})^\times \cong C(M)$ satisfies the smoothness condition. Also, by Theorem 3.24 $\mathbf{K}^{-1} > 0$. In other words $c_1^B(M)$ has a positive representative. By the transverse Calabi-Yau theorem [28] there is a transversal deformation $(M, \tilde{g}, \tilde{\Phi}, \tilde{\xi}, \tilde{\eta})$ with $\tilde{\eta} = \eta + 2d^c\phi$, for $\phi \in C_B^\infty(M)$, with $\text{Ric}_{\tilde{g}}^T$ positive. Then by (2.3) a homothetic deformation $\tilde{g}_a = a^2\tilde{\eta} \otimes \tilde{\eta} + a\tilde{g}^T$ for small enough $a \in \mathbb{R}_+$ has $\text{Ric}_{\tilde{g}_a}$ positive on M .

Considering $\pi : M \rightarrow X$ as an S^1 V-bundle over X , it lifts to a genuine fiber bundle over $B(X) = M \times_{S^1} E(S^1)$, $\tilde{\pi} : \tilde{M} \rightarrow B(X)$. Here $E(S^1)$ is the universal S^1 -principal bundle, and $B(X)$ is the orbifold classifying space (cf. [6]). Since M is smooth, we have a homotopy equivalence $\tilde{M} \simeq M$. And since $\pi_1^{org}(X) = e$, the first few terms of the Gysin sequence give

$$\mathbb{Z} \xrightarrow{\cup e} H_{orb}^2(X, \mathbb{Z}) \xrightarrow{\pi^*} H^2(M, \mathbb{Z}) \rightarrow 0,$$

where $e = c_1^{orb}(\mathbf{L}) \in H_{orb}^2(X, \mathbb{Z})$. We have $w_2(M) \equiv c_1(D) \bmod 2 \equiv \pi^*(c_1^{orb}(X)) \bmod 2$, which is zero precisely when $x = c_1^{orb}(X)$ is divisible by 2 in $H_{orb}^2(X, \mathbb{Z})/\mathbb{Z}(e)$. If this is the case then there is a $u \in H_{orb}^2(X, \mathbb{Z})$ with $2u = x + al$, with $a \in \mathbb{Z}$ odd. It is easy to see that $u \in H_{orb}^2(X, \mathbb{Z})$ is represented by a holomorphic line V-bundle, since $2u$ is. By Proposition 3.12, this V-bundle is $\mathbf{L}_u$ for some $u \in \text{SF}(\Delta^*)$, and we have an equation

$$(7.1) \quad 2u = -k + al + f, \quad \text{where } f \in M.$$

Evaluating (7.1) on σ_0 gives $2u(\sigma_0) = -1 + f(\sigma_0)$, and on $\sigma_2 = -\sigma_0$ gives $2u(\sigma_2) = -(1 + a) - f(\sigma_0)$. The first equation implies $f(\sigma_0)$ is odd, the second that $f(\sigma_0)$ is even. Thus $w_2(M) \neq 0$. Theorem 6.2 then completes the proof. $\square$

One can use arguments of M. Demazure (cf. [34] §3.4) as in the smooth case to show that $\text{Aut}^o(X)$ is not reductive for these examples. The fan $\Delta_{k,p}^*$ has two roots α_1, α_2 and $\alpha_2 \neq -\alpha_1$ as the Hirzebruch surface F_1 . The Lie algebra of Hamiltonian holomorphic vector fields of M is not reductive. By the proof of the Lichnerowicz theorem in the Sasakian case [10] the Sasakian structure can not be transversally deformed to constant scalar curvature.

Let $N_i, \bar{N}_i \subset \mathcal{S}, i = 1, \dots, k+2$ be the Sasakian submanifolds of the 3-Sasakian manifold $\mathcal{S}$ as discussed after Theorem 6.4. Suppose $w_2(\mathcal{M}) \neq 0$. So $N_i, \bar{N}_i$ are principle S^1 subbundles of $\mathbf{L}^{-1}$ restricted to toric surfaces $D, \bar{D}$ with fans as in Theorem 5.19. The augmented fan Δ^* of D has elements $\sigma_1 = \rho_1, \dots, \sigma_i = \rho_i, \sigma_{i+1} = -\rho_i + \rho_{i+1}, \sigma_{i+2} = \rho_{n+i+1}, \dots, \sigma_{k+3} = \rho_{2n}, n = k+2$, and $\mathbf{L}|_D$ is the line bundle associated to $l \in \text{SF}(\Delta^*)$ with

$$l(\sigma_j) := \begin{cases} 0 & \text{if } j = i+1, \\ -1 & \text{otherwise.} \end{cases}$$

The same argument in the above proposition shows that N_i is diffeomorphic to $X_\infty \# (k-1)M_\infty$.

8. HIGHER DIMENSIONAL EXAMPLES

In this section we employ the join construction of C. Boyer and K. Galicki [9, 6] to construct higher dimensional examples. Let $(M_i, g_i, \Phi_i, \xi_i, \eta_i), i = 1, 2$ be quasi-regular Sasakian manifolds of dimensions $2m_i + 1, i = 1, 2$. Make homothetic deformations of the Sasakian structures so that the S^1 -actions generated by $\xi_i, i = 1, 2$ have period 1. Then the transverse Kähler forms $\omega_i^T, i = 1, 2$ descend to forms $\omega_i, i = 1, 2$ on the leaf spaces $\mathcal{Z}_i, i = 1, 2$ with $[\omega_i] \in H_{orb}^2(\mathcal{Z}_i, \mathbb{Z}), i = 1, 2$. Then for a pair of positive integers (k_1, k_2) we have a kähler form $k_1\omega_1 + k_2\omega_2$ on $\mathcal{Z}_1 \times \mathcal{Z}_2$ with $[k_1\omega_1 + k_2\omega_2] \in H_{orb}^2(\mathcal{Z}_1 \times \mathcal{Z}_2, \mathbb{Z})$. By Proposition 2.4 there is an S^1 V-bundle, denoted $M_1 \star_{k_1, k_2} M_2$, with a homothetic family of Sasakian structures. We may assume that $\gcd(k_1, k_2) = 1$; for if $(k_1, k_2) = (lk'_1, lk'_2)$, $M_1 \star_{k_1, k_2} M_2 = (M_1 \star_{k'_1, k'_2} M_2)/\mathbb{Z}_l$. Note that

$$M_1 \star_{k_1, k_2} M_2 = (M_1 \times M_2)/S^1(k_1, k_2),$$

where S^1 acts by $(x, y) \rightarrow (e^{ik_2\theta}x, e^{-ik_1\theta}y)$.

In general the join $M_1 \star_{k_1, k_2} M_2$ is an orbifold of dimension $2(m_1 + m_2) + 1$. But a simple condition exists that implies smoothness. Let $v_i = \text{Ord}(M_i), i = 1, 2$ denote the lcm of the orders of the leaf holonomy groups of the $\mathcal{F}_{\xi_i}$.

Proposition 8.1 ([6, 9]). *For each pair (k_1, k_2) of relatively prime positive integers, $M_1 \star_{k_1, k_2} M_2$ is a smooth quasi-regular Sasakian manifold if, and only if, $\gcd(v_1k_2, v_2k_1) = 1$.*

Note that if M_1 and M_2 are positive, that is, $\text{Ricci}_{g_i}^T, i = 1, 2$ are positive, then $M_1 \star_{k_1, k_2} M_2$ is positive. Suppose M_1 and M_2 are Sasakian Einstein. We have $\text{Ind}(\mathcal{Z}_1 \times \mathcal{Z}_2) = \gcd(\text{Ind}(\mathcal{Z}_1), \text{Ind}(\mathcal{Z}_2))$. So we define the *relative indices* of M_1 and

M_2 to be

$$(8.1) \quad l_i = \frac{\text{Ind}(M_i)}{\gcd(\text{Ind}(M_1), \text{Ind}(M_2))}, \quad \text{for } i = 1, 2.$$

Then the homothetic family of Sasakian structures on the join $M_1 \star_{l_1, l_2} M_2$ has a Sasakian Einstein structure with transverse metric

$$g^T = \frac{(m_1 + 1)g_1^T + (m_2 + 1)g_2^T}{m_1 + m_2 + 1}.$$

Let $M_k \cong \#k(S^2 \times S^3)$ be one of the Sasaki-Einstein manifolds constructed in section 6. Then $\text{Ind}(M_k) = 1$ or 2 . Consider S^{2m+1} with its standard Sasakian structure. Then $\text{Ind}(S^{2m+1}) = m + 1$. Then the relative indices of S^{4j+3} and M_k are both 1. So Proposition 8.1 implies that $M_k \star S^{4j+3}$, $j \geq 0$, is a Sasaki-Einstein $(4j + 7)$ -manifold for all examples M_k . One can iterate this procedure; for example

$$M_k \star \overbrace{S^3 \star \cdots \star S^3}^{p \text{ times}},$$

is a $5 + 2p$ dimensional Sasaki-Einstein manifold. By making repeated joins to the examples in Corollary 6.6 we obtain the following.

Proposition 8.2. *For every possible dimension $n = 2m + 1 \geq 5$ there are infinitely many toric quasi-regular Sasaki-Einstein manifolds with arbitrarily high second Betti number.*

More precisely, in dimension 5 we have examples of every odd b_2 , and infinitely many examples for each odd $b_2 \geq 3$.

In dimension $n = 2m + 1 \geq 7$ for m odd we have examples of every even b_2 , and infinitely many examples for each even $b_2 \geq 4$.

In dimension $n = 2m + 1 \geq 9$ for m even we have examples of every odd b_2 , and infinitely many examples for each odd $b_2 \geq 5$.

Example. Consider the 7-dimensional Sasaki-Einstein manifolds $S^3 \star M_k$. The Gysin sequence of $\pi : S^3 \star M_k \rightarrow \mathbb{C}P^1 \times X$ determines the cohomology in $\mathbb{Q}$ -coefficients. The Leray spectral sequence of the fiber bundle $\varpi : S^3 \star M_k \rightarrow \mathbb{C}P^1$ with fiber M_k can be used to show the following.

p	0	1	2	3	4	5	6	7
$H^p(S^3 \star M_k, \mathbb{Z})$	$\mathbb{Z}$	0	$\mathbb{Z}^{k+1}$	0	T	$\mathbb{Z}^{k+1}$	0	$\mathbb{Z}$

Here T is a torsion group. This is quite similar to the cohomology of S_Ω in Theorem 5.14. $\diamond$

Let $N = N_{k,p}$ denote the positive Ricci curvature Sasakian 5-manifold of Proposition 7.1 diffeomorphic to $X_\infty \# (k - 1)M_\infty$. Then we see by Proposition 8.1 that for any regular Sasaki-Einstein manifold M , $N \star_{1,l} M$ is smooth for any $l \geq 1$. Furthermore, we may transversally deform it to a positive Ricci curvature structure.

Let $\pi_i : \mathbf{L}_i \rightarrow Z_i, i = 1, 2$ be the holomorphic V-bundles whose S^1 -principal bundles are N and M respectively. Then $N \star_{1,l} M$ is the principal S^1 V-bundle of $\pi : \mathbf{L}_1 \otimes \mathbf{L}_2^l$, and as in the proof of Proposition 7.1 we have

$$\mathbb{Z} \xrightarrow{\cup e} H_{orb}^2(Z_1 \times Z_2, \mathbb{Z}) \xrightarrow{\pi^*} H^2(N \star_{1,l} M, \mathbb{Z}) \rightarrow 0,$$

where $e = c_1(\mathbf{L}_1) + lc_1(\mathbf{L}_2)$. We have $w_2(N \star_{1,l} M) \equiv \pi^*(c_1(Z_1) + c_1(Z_2)) \pmod{2}$, and $w_2(N \star_{1,l} M) = 0$ if, and only if, $\pi^*(c_1(Z_1) + c_1(Z_2))$ is divisible by 2 in $H^2(N \star_{1,l} M, \mathbb{Z})$. If this is the case then there is an $u \in H^2_{orb}(Z_1 \times Z_2, \mathbb{Z})$ with

$$2u = c_1(Z_1) + c_1(Z_2) + s(c_1(\mathbf{L}_1) + lc_1(\mathbf{L}_2)), \quad \text{for } s \in \mathbb{Z}.$$

Let $\iota : Z_1 \rightarrow Z_1 \times Z_2$ be the inclusion $\iota(x) = (x, y)$ with $y \in Z_2$ a smooth point. Then $2\iota^*u = c_1(Z_1) + sc_1(\mathbf{L}_1)$ which contradicts Proposition 7.1. Thus $w_2(N \star_{1,l} M) \neq 0$. By taking joins of the $N_{k,p}$, $k \geq 1, p \geq 0$ with spheres as above we obtain the following.

Proposition 8.3. *For every dimension $n = 2m + 1 \geq 5$ there exist infinitely many toric quasi-regular positive Ricci curvature Sasakian manifolds of each $b_2 \geq 2$, and of each $b_2 \geq 1$ in dimension 5. These examples are simply connected and have $w_2 \neq 0$. Therefore, they do not admit a Sasaki-Einstein structure.*

REFERENCES

- [1] M. F. Atiyah, N. J. Hitchin, and I.M. Singer. Self-duality in four-dimensional Riemannian geometry. *Proc. Roy. Soc. London A*, 362:425–461, 1978.
- [2] W. L. Baily. On the imbedding of V-manifolds in projective space. *Amer. J. Math.*, 79:403–430, 1957.
- [3] D. Barden. Simply connected five-manifolds. *Ann. of Math.*, 82(2):365–385, 1965.
- [4] R. Bielawski. Complete hyper-Kähler $4n$ -manifolds with a local tri-Hamiltonian $\mathbf{R}^n$ -action. *Math. Ann.*, 314(3):505–528, 1999.
- [5] C. P. Boyer and K. Galicki. 3-Sasakian manifolds. In *Surveys in Differential Geometry: Einstein Manifolds*, Surv. Differ. Geom., VI, pages 123–184. Int. Press, Boston, MA, 1999.
- [6] C. P. Boyer and K. Galicki. On Sasakian-Einstein geometry. *Internat. J. Math.*, 11(7):873–909, 2000.
- [7] C. P. Boyer, K. Galicki, and B. M. Mann. Hypercomplex structures from 3-Sasakian structures. *J. Reine Angew. Math.*, 501:115–141, 1998.
- [8] C. P. Boyer, K. Galicki, B. M. Mann, and E. G. Rees. Compact 3-Sasakian 7 manifolds with arbitrary second Betti number. *Invent. Math.*, 131(2):321–344, 1998.
- [9] C. P. Boyer, K. Galicki, and L. Ornea. Constructions in Sasakian geometry. *Math. Z.*, 257(4):907–924, 2007.
- [10] C. P. Boyer, K. Galicki, and S. Simanca. Canonical Sasakian metrics. *Comm. Math. Phys.*, 279(3):705–733, 2008.
- [11] D. Calderbank, L. David, and P. Gauduchon. The Guillemin formula and Kähler metrics on toric symplectic manifolds. *J. Symplectic Geom.*, 1(4):767–784, 2003.
- [12] D. Calderbank and M. Singer. Toric self-dual Einstein metrics on compact orbifolds. *Duke Math. J.*, 133(2):237–258, 2006.
- [13] T. Delzant. Hamiltoniens périodiques et image convexe de l’application moment. *Bull. Soc. Math. France*, 116:315–339, 1988.
- [14] J. H. Eschenburg. New examples of manifolds with strictly positive curvature. *Invent. Math.*, 66:469–480, 1982.
- [15] Th. Friedrich and H. Kurke. Compact four-dimensional self-dual Einstein manifolds with positive scalar curvature. *Math. Nachr.*, 106:271–299, 1982.
- [16] W. Fulton. *Introduction to Toric Varieties*. Number 131 in Annals of Mathematics Studies. Princeton University Press, Princeton, New Jersey, 1993.
- [17] A. Futaki. An obstruction to the existence of Kähler-Einstein metrics. *Invent. Math.*, 73:437–443, 1983.
- [18] A. Futaki, H. Ono, and G. Wang. Transverse Kähler geometry of Sasaki manifolds and toric Sasaki-Einstein manifolds. preprint arXiv:math.DG/0607586 v.5, January 2007.
- [19] K. Galicki and H. B. Lawson. Quaternionic reduction and quaternionic orbifolds. *Math. Ann.*, 282:1–21, 1988.
- [20] A. Grothendieck. *Sur les faisceaux algébriques et les faisceaux analytiques cohérents*, volume 16 of *Séminaire Henri Cartan; 9e année: 1956/57. Quelques questions de topologie, Exposé no. 2*. Secrétariat mathématique, Paris, 1958.

- [21] A. Grothendieck. Géométrie algébriques et géométrie analytique (notes by Mme. M. Raymond), SGA 1960/61, Exp. XII. In *Revêtements étales et groupe fondamental*, volume 224 of *Lecture Notes in Math.* Springer-Verlag, Heidelberg, 1971.
- [22] V. Guillemin. Kähler structures on toric varieties. *J. Diff. Geo.*, 40:285–309, 1994.
- [23] V. Guillemin. *Moment Maps and Combinatorial Invariants of Hamiltonian T^n -spaces*, volume 122 of *Progress in Mathematics*. Birkhäuser, 1994.
- [24] A. Haefliger. Groupoïdes d’holonomie et classifiants. *Astérisque*, (116):70–97, 1984. Transversal structure of foliations (Toulouse, 1982).
- [25] A. Haefliger and E. Salem. Actions of tori on orbifolds. *Ann. Global Anal. Geom.*, 9(1):37–59, 1991.
- [26] R. Hepworth. The topology of certain 3-Sasakian 7-manifolds. *Math. Ann.*, 339(4):733–755, 2007.
- [27] N. J. Hitchin. Kählerian twistor spaces. *Proc. London Math. Soc. (3)*, 43(1):133–150, 1981.
- [28] A. El Kacimi-Alaoui. Opérateurs transversalement elliptiques sur un feuilletage riemannien et applications. *Compositio Mathematica*, 79:57–106, 1990.
- [29] E. Lerman and S. Tolman. Hamiltonian torus actions on symplectic orbifolds and symplectic varieties. *Trans. Amer. Math. Soc.*, 349:4201–4230, 1997.
- [30] T. Mabuchi. Einstein-Kähler forms, Futaki invariants and convex geometry on toric Fano varieties. *Osaka J. Math.*, 24:705–737, 1987.
- [31] D. Martelli, J. Sparks, and S.-T. Yau. The geometric dual of a -maximisation for toric Sasaki-Einstein manifolds. *Comm. Math. Phys.*, 268(1):39–65, 2006.
- [32] D. Martelli, J. Sparks, and S.-T. Yau. Sasaki-Einstein manifolds and volume minimisation. *Comm. Math. Phys.*, 280(3):611–673, 2008.
- [33] T. Oda. *Lectures on Torus Embeddings and Applications (Based on Joint Work with Katsuya Miyake)*, volume 58 of *Tata Inst. Inst. of Fund. Research*. Springer, Berlin, Heidelberg, New York, 1978.
- [34] T. Oda. *Convex Bodies in Algebraic Geometry*, volume 15 of *Ergebnisse der Math. u. ihrer Grenz. Geb. 3 Folge*. Springer, 1985.
- [35] H. S. Oh. Toral actions on 5-manifolds. *Trans. Amer. Math.*, 278(1):233–252, 1983.
- [36] R. Penrose. Nonlinear gravitons and curved twistor theory. *General Relativity and Gravitation*, 7(1):31–52, 1976. The riddle of gravitation—on the occasion of the 60th birthday of Peter G. Bergmann (Proc. Conf., Syracuse Univ., Syracuse, N. Y., 1975).
- [37] S. Salamon. Quaternionic Kähler manifolds. *Invent. Math.*, 67:143–171, 1982.
- [38] S. Smale. On the structure of 5-manifolds. *Annals of Math.*, 75(1):38–46, 1962.
- [39] G. Tian and X. Zhu. A new holomorphic invariant and uniqueness of Kähler-Ricci solitons. *Comment. Math. Helv.*, 77:297–325, 2002.
- [40] C. van Coevering. Toric surfaces and Sasakian-Einstein 5-manifolds. preprint arXiv:math.DG/0607721 v1, July 2006.
- [41] X. J. Wang and X. Zhu. Kähler-Ricci solitons on toric manifolds with positive first Chern class. *Advances in Math.*, 188:87–103, 2004.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, 77 MASSACHUSETTS AVENUE, CAMBRIDGE, MA 02139-4307

E-mail address: craig@math.mit.edu